

ABALOPARATIDE

Products Affected

- Tymlos

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	ONE OF THE FOLLOWING: (1) HIGH RISK FOR FRACTURES DEFINED AS ONE OF THE FOLLOWING: HISTORY OF OSTEOPOROTIC (I.E., FRAGILITY, LOW TRAUMA) FRACTURE(S). 2 OR MORE RISK FACTORS FOR FRACTURE (E.G., HISTORY OF MULTIPLE RECENT LOW TRAUMA FRACTURES, BMD T-SCORE LESS THAN OR EQUAL TO -2.5, CORTICOSTEROID USE, OR USE OF GNRH ANALOGS SUCH AS NAFARELIN, ETC.). NO PRIOR TREATMENT FOR OSTEOPOROSIS AND FRAX SCORE OF AT LEAST 20% FOR ANY MAJOR FRACTURE OR OF AT LEAST 3% FOR HIP FRACTURE. (2) UNABLE TO USE ORAL THERAPY (I.E., UPPER GASTROINTESTINAL PROBLEMS UNABLE TO TOLERATE ORAL MEDICATION, LOWER GASTROINTESTINAL PROBLEMS UNABLE TO ABSORB ORAL MEDICATIONS, TROUBLE REMEMBERING TO TAKE ORAL MEDICATIONS OR COORDINATING AN ORAL BISPHOSPHONATE WITH OTHER ORAL MEDICATIONS OR THEIR DAILY ROUTINE). (3) ADEQUATE TRIAL OF, INTOLERANCE TO, OR A CONTRAINDICATION TO BISPHOSPHONATES (E.G., ALENDRONATE, RISEDRONATE, IBANDRONATE).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	

ABATACEPT IV

Products Affected

- Orenzia (with maltose)

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	RENEWAL: PHYSICIAN ATTESTATION OF IMPROVEMENT.
Age Restrictions	
Prescriber Restrictions	PRESCRIBED BY OR GIVEN IN CONSULTATION WITH A RHEUMATOLOGIST
Coverage Duration	INITIAL: RA: 6 MONTHS. JIA: 4 MONTHS. RENEWAL: 12 MONTHS
Other Criteria	INITIAL: RHEUMATOID ARTHRITIS (RA): PREVIOUS TRIAL OF HUMIRA AND ONE DMARD (DISEASE-MODIFYING ANTIRHEUMATIC DRUG) SUCH AS METHOTREXATE, LEFLUNOMIDE, HYDROXYCHLOROQUINE, OR SULFASALAZINE. JUVENILE IDIOPATHIC ARTHRITIS (JIA): PREVIOUS TRIAL OF HUMIRA AND ONE DMARD (DISEASE-MODIFYING ANTIRHEUMATIC DRUG) SUCH AS METHOTREXATE, LEFLUNOMIDE, HYDROXYCHLOROQUINE, OR SULFASALAZINE.

ABATACEPT SQ

Products Affected

- Orenzia
- Orenzia ClickJect

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	RENEWAL: PHYSICIAN ATTESTATION OF IMPROVEMENT.
Age Restrictions	
Prescriber Restrictions	PRESCRIBED BY OR GIVEN IN CONSULTATION WITH A RHEUMATOLOGIST.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.
Other Criteria	INITIAL: RHEUMATOID ARTHRITIS (RA): PREVIOUS TRIAL OF HUMIRA AND ONE DMARD (DISEASE-MODIFYING ANTIRHEUMATIC DRUG) SUCH AS METHOTREXATE, LEFLUNOMIDE, HYDROXYCHLOROQUINE, OR SULFASALAZINE. JUVENILE IDIOPATHIC ARTHRITIS (JIA): PREVIOUS TRIAL OF HUMIRA AND ONE DMARD (DISEASE-MODIFYING ANTIRHEUMATIC DRUG) SUCH AS METHOTREXATE, LEFLUNOMIDE, HYDROXYCHLOROQUINE, OR SULFASALAZINE.

ABIRATERONE

Products Affected

- Zytiga

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	

ADALIMUMAB

Products Affected

- Humira
- Humira Pediatric Crohn's Start
- Humira Pen
- Humira Pen Crohn's-UC-HS Start
- Humira Pen Psoriasis-Uveitis

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	INITIAL: POLYARTICULAR JUVENILE IDIOPATHIC ARTHRITIS: CURRENT WEIGHT. PLAQUE PSORIASIS: MODERATE TO SEVERE PLAQUE PSORIASIS INVOLVING GREATER THAN OR EQUAL TO 5% BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, OR GENITAL AREA. RENEWAL: PHYSICIAN ATTESTATION OF IMPROVEMENT.
Age Restrictions	
Prescriber Restrictions	RHEUMATOID ARTHRITIS, POLYARTICULAR JUVENILE IDIOPATHIC ARTHRITIS, ANKYLOSING SPONDYLITIS: PRESCRIBED BY OR GIVEN IN CONSULTATION WITH A RHEUMATOLOGIST. PSORIATIC ARTHRITIS: PRESCRIBED BY OR GIVEN IN CONSULTATION WITH A DERMATOLOGIST OR RHEUMATOLOGIST. PSORIASIS: PRESCRIBED BY OR GIVEN IN CONSULTATION WITH A DERMATOLOGIST. CROHN'S DISEASE/ULCERATIVE COLITIS: PRESCRIBED BY OR GIVEN IN CONSULTATION WITH A GASTROENTEROLOGIST.
Coverage Duration	INITIAL:RA:6 MO PSA/AS:4 MO PJIA:5 MO PSO/CD/UC/HS:3 MO UVEITIS:6 MO RENEWAL:12 MO ALL INDICATIONS

PA Criteria	Criteria Details
Other Criteria	<p>INITIAL: RHEUMATOID ARTHRITIS (RA): PREVIOUS TRIAL OF ONE DMARD (DISEASE-MODIFYING ANTIRHEUMATIC DRUG) SUCH AS METHOTREXATE, LEFLUNOMIDE, HYDROXYCHLOROQUINE, OR SULFASALAZINE. POLYARTICULAR JUVENILE IDIOPATHIC ARTHRITIS (PJIA): PREVIOUS TRIAL OF ONE DMARD (DISEASE-MODIFYING ANTIRHEUMATIC DRUG) SUCH AS METHOTREXATE, LEFLUNOMIDE, HYDROXYCHLOROQUINE, OR SULFASALAZINE. PSORIATIC ARTHRITIS (PSA): PREVIOUS TRIAL OF ONE DMARD (DISEASE-MODIFYING ANTIRHEUMATIC DRUG) SUCH AS METHOTREXATE, LEFLUNOMIDE, HYDROXYCHLOROQUINE, OR SULFASALAZINE. ANKYLOSING SPONDYLITIS: TRIAL OF FORMULARY AGENTS NOT REQUIRED. PLAQUE PSORIASIS (PSO): PREVIOUS TRIAL OF ONE OF THE FOLLOWING CONVENTIONAL THERAPIES SUCH AS PUVA (PHOTOTHERAPY ULTRAVIOLET LIGHT A), UVB (ULTRAVIOLET LIGHT B), TOPICAL CORTICOSTEROIDS, CALCIPOTRIENE, ACITRETIN, METHOTREXATE, OR CYCLOSPORINE. CROHN'S DISEASE (CD): PREVIOUS TRIAL OF ONE CONVENTIONAL AGENT SUCH AS A CORTICOSTEROID (I.E., BUDESONIDE, METHYLPREDNISOLONE), AZATHIOPRINE, MERCAPTOPURINE, METHOTREXATE, OR MESALAMINE. ULCERATIVE COLITIS (UC): PREVIOUS TRIAL OF ONE CONVENTIONAL AGENT SUCH AS A CORTICOSTEROID (I.E., BUDESONIDE, METHYLPREDNISOLONE), AZATHIOPRINE, MERCAPTOPURINE, METHOTREXATE, OR MESALAMINE.</p>

AFATINIB DIMALEATE

Products Affected

- Gilotrif

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	

ALECTINIB

Products Affected

- Alecensa

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	

ALEMTUZUMAB - LEMTRADA

Products Affected

- Lemtrada

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	INITIAL: 1 MONTH. RENEWAL: 12 MONTHS.
Other Criteria	TRIAL WITH TWO OF THE FOLLOWING AGENTS FOR MULTIPLE SCLEROSIS: AUBAGIO, AVONEX, GILENYA, PLEGRIDY, REBIF, TECFIDERA, OR GLATIRAMER. RENEWAL REQUESTS FOR ALEMTUZUMAB REQUIRE THAT AT LEAST 12 MONTHS HAVE ELAPSED SINCE RECEIVING THE FIRST COURSE OF LEMTRADA. PATIENTS ARE LIMITED TO TWO COURSES OF THERAPY WITH LEMTRADA WITHIN A LIFETIME.

ALIROCUMAB

Products Affected

- Praluent Pen

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	18 YEARS OF AGE AND OLDER.
Prescriber Restrictions	CARDIOLOGIST, ENDOCRINOLOGIST OR LIPIDOLOGIST
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS

PA Criteria	Criteria Details
Other Criteria	<p>MUST HAVE A LDL CHOLESTEROL LEVEL GREATER THAN 100MG/DL WHILE ON MAXIMAL DRUG TREATMENT FOR THE PAST 2 MONTHS AND ONE OF THE FOLLOWING DIAGNOSES: (1) HETEROZYGOUS FAMILIAL HYPERCHOLESTEROLEMIA (HEFH) DETERMINED BY SIMON BROOME DIAGNOSTIC CRITERIA FOR HEFH OR A SCORE OF 6 OR GREATER ON THE DUTCH LIPID NETWORK CRITERIA FOR HEFH OR (2) HISTORY OF ATHEROSCLEROTIC CARDIOVASCULAR DISEASE (ASCVD) AS DOCUMENTED BY PHYSICIAN ATTESTATION. PATIENT MUST NOT HAVE CONCURRENT USE OF REPATHA OR OTHER PCSK9 AGENT. INITIAL THERAPY: FOR STATIN TOLERANT PATIENTS: MUST HAVE TAKEN ATORVASTATIN OR ROSUVASTATIN FOR THE PAST 2 MONTHS. FOR STATIN INTOLERANT PATIENTS: DOCUMENTATION OF STATIN INTOLERANCE BY ONE OF THE FOLLOWING: (1) PHYSICIAN ATTESTATION, (2) PATIENT HAS TRIED ROSUVASTATIN, ATORVASTATIN, OR STATIN THERAPY AT ANY DOSE AND HAS EXPERIENCED SKELETAL-MUSCLE RELATED SYMPTOMS (E.G., MYOPATHY). PATIENTS WITH CONTRAINDICATIONS TO STATINS INCLUDING ACTIVE DECOMPENSATED LIVER DISEASE, NURSING FEMALE, PREGNANCY OR PLANS TO BECOME PREGNANT OR HYPERSENSITIVITY REACTIONS WILL BE APPROVED FOR PRALUENT THERAPY WITHOUT REQUIREMENT OF DOCUMENTATION OF STATIN INTOLERANCE. RENEWAL CRITERIA: RECEIVING PRIOR PRALUENT THERAPY FOR THE PAST 6 MONTHS AND NO CLAIMS FOR REPATHA, JUXTAPID, OR KYNAMRO SINCE PRALUENT APPROVAL.</p>

ANAKINRA

Products Affected

- Kineret

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	RENEWAL: PHYSICIAN ATTESTATION OF IMPROVEMENT.
Age Restrictions	
Prescriber Restrictions	RHEUMATOID ARTHRITIS: PRESCRIBED BY OR GIVEN IN CONSULTATION WITH A RHEUMATOLOGIST.
Coverage Duration	INITIAL: RA: 6 MONTHS NOMID/CAPS: 12 MONTHS. RENEWAL: 12 MONTHS.
Other Criteria	INITIAL: RHEUMATOID ARTHRITIS (RA): PREVIOUS TRIAL OF HUMIRA FOLLOWED BY ONE OF THE FOLLOWING PREFERRED AGENTS: ORENCIA, XELJANZ, CIMZIA, OR ACTEMRA.

APREMILAST

Products Affected

- Otezla
- Otezla Starter

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	INITIAL: MODERATE TO SEVERE PLAQUE PSORIASIS INVOLVING GREATER THAN OR EQUAL TO 5% OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, OR GENITAL AREA. RENEWAL: PHYSICIAN ATTESTATION OF IMPROVEMENT.
Age Restrictions	
Prescriber Restrictions	PSORIATIC ARTHRITIS: PRESCRIBED BY OR GIVEN IN CONSULTATION WITH A DERMATOLOGIST OR RHEUMATOLOGIST. PSORIASIS: PRESCRIBED BY OR GIVEN IN CONSULTATION WITH A DERMATOLOGIST
Coverage Duration	INITIAL: PSORIATIC ARTHRITIS: 4 MONTHS. PSORIASIS: 5 MONTHS. RENEWAL: 12 MONTHS
Other Criteria	INITIAL: PSORIATIC ARTHRITIS (PSA): PREVIOUS TRIAL OF HUMIRA AND ONE DMARD (DISEASE-MODIFYING ANTIRHEUMATIC DRUG) SUCH AS METHOTREXATE, LEFLUNOMIDE, HYDROXYCHLOROQUINE, OR SULFASALAZINE. PLAQUE PSORIASIS (PSO): PREVIOUS TRIAL OF HUMIRA AND ONE CONVENTIONAL THERAPY SUCH AS PUVA (PHOTOTHERAPY ULTRAVIOLET LIGHT A), UVB (ULTRAVIOLET LIGHT B), TOPICAL CORTICOSTEROIDS, CALCIPOTRIENE, ACITRETIN, METHOTREXATE, OR CYCLOSPORINE.

ASFOTASE

Products Affected

- Strensiq

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	NON-SPECIFIC ALKALINE PHOSPHATASE (TNSALP) (ALPL) GENE MUTATION, SERUM ALKALINE PHOSPHATASE (ALP) LEVEL, SERUM PYRIDOXAL-5'-PHOSPHATE (PLP) LEVELS, URINE PHOSPHOETHANOLAMINE (PEA) LEVEL, RADIOGRAPHIC EVIDENCE OF HYPOPHOSPHATASIA (HPP)
Age Restrictions	PERINATAL/INFANTILE-ONSET HYPOPHOSPHATASIA (HPP): 6 MONTHS OF AGE OR YOUNGER AT HYPOPHOSPHATASIA (HPP) ONSET. JUVENILE-ONSET HYPOPHOSPHATASIA (HPP): 18 YEARS OF AGE OR YOUNGER AT HYPOPHOSPHATASIA (HPP) ONSET.
Prescriber Restrictions	PRESCRIBED BY OR GIVEN IN CONSULTATION WITH AN ENDOCRINOLOGIST, A GENETICIST, OR A METABOLIC SPECIALIST.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.

PA Criteria	Criteria Details
Other Criteria	<p>INITIAL: FOR PATIENTS WITH PERINATAL/INFANTILE-ONSET HYPOPHOSPHATASIA (HPP), ALL OF THE FOLLOWING CRITERIA MUST BE MET: POSITIVE FOR A TISSUE NON-SPECIFIC ALKALINE PHOSPHATASE (TNSALP) (ALPL) GENE MUTATION AS CONFIRMED BY GENETIC TESTING OR MEETS AT LEAST TWO OF THE FOLLOWING CRITERIA: 1.) SERUM ALKALINE PHOSPHATASE (ALP) LEVEL BELOW THAT OF NORMAL RANGE FOR PATIENT AGE 2.) SERUM PYRIDOXAL-5'-PHOSPHATE (PLP) LEVELS ELEVATED AND PATIENT HAS NOT RECEIVED VITAMIN B6 SUPPLEMENTATION IN THE PREVIOUS WEEK 3.) URINE PHOSPHOETHANOLAMINE (PEA) LEVEL ABOVE THAT OF NORMAL RANGE FOR PATIENT AGE 4.) RADIOGRAPHIC EVIDENCE OF HYPOPHOSPHATASIA (HPP) (E.G., FLARED AND FRAYED METAPHYSES, OSTEOPENIA, WIDENED GROWTH PLATES, AREAS OF RADIOLUCENCY OR SCLEROSIS) 5.) PRESENCE OF TWO OR MORE OF THE FOLLOWING: RACHITIC CHEST DEFORMITY, CRANIOSYNOSTOSIS (PREMATURE CLOSURE OF SKULL BONES), DELAY IN SKELETAL GROWTH RESULTING IN DELAY OF MOTOR DEVELOPMENT, HISTORY OF VITAMIN B6 DEPENDENT SEIZURES, NEPHROCALCINOSIS, OR HISTORY OF ELEVATED SERUM CALCIUM. HISTORY OR PRESENCE OF NON-TRAUMATIC POSTNATAL FRACTURE AND DELAYED FRACTURE HEALING. FOR PATIENTS WITH JUVENILE-ONSET HYPOPHOSPHATASIA (HPP), ALL OF THE FOLLOWING CRITERIA MUST BE MET: POSITIVE FOR A TISSUE NON-SPECIFIC ALKALINE PHOSPHATASE (TNSALP) (ALPL) GENE MUTATION AS CONFIRMED BY GENETIC TESTING OR MEETS AT LEAST TWO OF THE FOLLOWING CRITERIA: 1.) SERUM ALKALINE PHOSPHATASE (ALP) LEVEL BELOW THAT OF NORMAL RANGE FOR PATIENT AGE 2.) SERUM PYRIDOXAL-5'-PHOSPHATE (PLP) LEVELS ELEVATED AND PATIENT HAS NOT RECEIVED VITAMIN B6 SUPPLEMENTATION IN THE PREVIOUS WEEK 3.)URINE PHOSPHOETHANOLAMINE (PEA) LEVEL ABOVE THAT OF NORMAL RANGE FOR PATIENT AGE 4.)RADIOGRAPHIC EVIDENCE OF HYPOPHOSPHATASIA (HPP) (E.G., FLARED AND FRAYED METAPHYSES, OSTEOPENIA, OSTEOMALACIA, WIDENED GROWTH PLATES, AREAS OF RADIOLUCENCY OR SCLEROSIS) 5.)PRESENCE OF TWO OR MORE OF THE FOLLOWING:RACHITIC DEFORMITIES (RACHITIC CHEST,</p>

PA Criteria	Criteria Details
	<p>BOWED LEGS, KNOCK-KNEES),PREMATURE LOSS OF PRIMARY TEETH PRIOR TO 5 YEARS OF AGE, DELAY IN SKELETAL GROWTH RESULTING IN DELAY OF MOTOR DEVELOPMENT, OR HISTORY OR PRESENCE OF NON-TRAUMATIC FRACTURES OR DELAYED FRACTURE HEALING. STRENSIQ WILL NOT BE APPROVED FOR THE FOLLOWING PATIENTS: PATIENTS CURRENTLY RECEIVING TREATMENT WITH A BISPHOSPHONATE [E.G., BONIVA (IBANDRONATE), FOSAMAX (ALENDRONATE), ACTONEL (RISEDRONATE)], PATIENTS WITH SERUM CALCIUM OR PHOSPHATE LEVELS BELOW THE NORMAL RANGE, PATIENTS WITH A TREATABLE FORM OF RICKETS. RENEWAL: PATIENT HAS EXPERIENCED AN IMPROVEMENT IN THE SKELETAL CHARACTERISTICS OF HYPOPHOSPHATASIA (HPP) (E.G., IMPROVEMENT OF THE IRREGULARITY OF THE PROVISIONAL ZONE OF CALCIFICATION, PHYSEAL WIDENING, METAPHYSEAL FLARING, RADIOLUCENCIES, PATCHY OSTEOSCLEROSIS, RATIO OF MID-DIAPHYSEAL CORTEX TO BONE THICKNESS, GRACILE BONES, BONE FORMATION AND FRACTURES.</p>

ASPARAGINASE

Products Affected

- Oncaspar

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	3 MONTHS
Other Criteria	

ATEZOLIZUMAB

Products Affected

- Tecentriq

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	

AVELUMAB

Products Affected

- Bavencio

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	

AXITINIB

Products Affected

- Inlyta oral tablet 1 mg, 5 mg

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	TRIAL OF AT LEAST ONE SYSTEMIC THERAPY FOR THE TREATMENT OF RCC SUCH AS NEXAVAR (SORAFENIB), TORISEL (TEMSIROLIMUS), SUTENT (SUNITINIB), VOTRIENT (PAZOPANIB), OR AVASTIN (BEVACIZUMAB) IN COMBINATION WITH INTERFERON.

BECAPLERMIN

Products Affected

- Regranex

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	NON-DIABETIC ULCERS, NEOPLASM AT APPLICATION SITE, PRESSURE OR VENOUS STASIS ULCERS AND ULCERS THAT DO NOT EXTEND THROUGH THE DERMIS.
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	VASCULAR SURGEON, PODIATRIST, ENDOCRINOLOGIST, PHYSICIAN PRACTICING IN A SPECIALTY WOUND CLINIC OR INFECTIOUS DISEASE SPECIALIST.
Coverage Duration	3 MONTHS
Other Criteria	

BEDAQUILINE FUMARATE

Products Affected

- Sirturo

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	18 YEARS OF AGE AND OLDER.
Prescriber Restrictions	
Coverage Duration	24 WEEKS
Other Criteria	SIRTURO USED IN COMBINATION WITH AT LEAST 3 OTHER ANTIBIOTICS FOR THE TREATMENT OF PULMONARY MULTI-DRUG RESISTANT TUBERCULOSIS.

BELIMUMAB

Products Affected

- Benlysta intravenous

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	AUTOANTIBODY POSITIVE LUPUS TEST.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS
Other Criteria	INITIAL: MEMBER IS CURRENTLY TAKING CORTICOSTEROIDS, ANTIMALARIALS, NSAIDS, OR IMMUNOSUPPRESSIVE AGENTS. NO APPROVAL FOR DIAGNOSIS OF SEVERE ACTIVE LUPUS NEPHRITIS, SEVERE CENTRAL NERVOUS SYSTEM LUPUS OR CONCURRENT USE OF BIOLOGIC AGENTS OR INTRAVENOUS CYCLOPHOSPHAMIDE. RENEWAL: PHYSICIAN ATTESTATION OF IMPROVEMENT.

BELINOSTAT

Products Affected

- Beleodaq

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	

BENDAMUSTINE

Products Affected

- Bendeka

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	

BEVACIZUMAB

Products Affected

- Avastin

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	

BEXAROTENE

Products Affected

- bexarotene
- Targretin topical

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	

BLINATUMOMAB

Products Affected

- Blincyto intravenous kit

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	INITIAL: 3 MONTHS. RENEWAL: 12 MONTHS
Other Criteria	INITIAL: DIAGNOSIS OF PHILADELPHIA CHROMOSOME-NEGATIVE (PH-) B-CELL PRECURSOR ACUTE LYMPHOBLASTIC LEUKEMIA IN A PATIENT WHO HAS PREVIOUSLY TRIED CHEMOTHERAPY BUT HAS RELAPSED OR IS REFRACTORY TO TREATMENT. INITIAL APPROVAL IS FOR 2 CYCLES, MAY APPROVE FOR 1 ADDITIONAL CYCLE DUE TO TREATMENT INTERRUPTION FOR DOSE MODIFICATION. RENEWAL IS APPROVED FOR PATIENTS WHO HAVE ACHIEVED COMPLETE REMISSION WITH OR WITHOUT PARTIAL HEMATOLOGICAL RECOVERY OF PERIPHERAL BLOOD COUNTS AFTER 2 CYCLES OF TREATMENT. RENEWAL IS NOT APPROVED FOR PATIENTS WHO RECEIVED AN ALLOGENEIC HEMATOPOIETIC STEM-CELL TRANSPLANT.

BORTEZOMIB

Products Affected

- Velcade

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	

BOSUTINIB

Products Affected

- Bosulif oral tablet 100 mg, 500 mg

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	CML: BCR-ABL MUTATIONAL ANALYSIS CONFIRMING THAT BOTH T315I AND V299L MUTATIONS ARE NOT PRESENT.

BOTULINUM NEUROTOXIN

Products Affected

- Botox injection recon soln 100 unit, 200 unit
- Myobloc intramuscular solution 10,000 unit/2 mL, 2,500 unit/0.5 mL, 5,000 unit/mL

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	COSMETIC DIAGNOSIS: WRINKLES.
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	MIGRAINE HEADACHE: TRIAL OF TWO OF THE FOLLOWING: BETA BLOCKERS, TRICYCLIC ANTIDEPRESSANTS, OR VALPROIC ACID. OVERACTIVE BLADDER: TRIAL OF OR CONTRAINDICATION TO THE USE OF ONE ANTICHOLINERGIC MEDICATION SUCH AS ORAL OXYBUTYNIN, ORAL OXYBUTYNIN ER, TOLTERODINE, TOLTERODINE ER, TOVIAZ, TROSPIUM, OR TROSPIUM ER. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION.

BRIGATINIB

Products Affected

- Alunbrig

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	

BRODALUMAB

Products Affected

- Siliq

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	INITIAL: MODERATE TO SEVERE PLAQUE PSORIASIS INVOLVING GREATER THAN OR EQUAL TO 5% OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, OR GENITAL AREA. RENEWAL: PHYSICIAN ATTESTATION OF IMPROVEMENT.
Age Restrictions	
Prescriber Restrictions	PRESCRIBED BY OR GIVEN IN CONSULTATION WITH A DERMATOLOGIST.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS
Other Criteria	INITIAL: PLAQUE PSORIASIS (PSO): PREVIOUS TRIAL OF HUMIRA FOLLOWED BY ONE OF THE FOLLOWING: COSENTYX OR OTEZLA. PATIENT HAS BEEN COUNSELED ON AND EXPRESSES UNDERSTANDING OF THE RISK OF SUICIDAL IDEATION AND BEHAVIOR. RENEWAL: PATIENT HAS NOT DEVELOPED OR REPORTED WORSENING DEPRESSIVE SYMPTOMS OR SUICIDAL IDEATION AND BEHAVIORS WHILE ON TREATMENT WITH SILIQ.

C1 ESTERASE INHIBITOR

Products Affected

- Cinryze
- Haegarda

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	HEMATOLOGIST, IMMUNOLOGIST
Coverage Duration	12 MONTHS
Other Criteria	

CABOZANTINIB

Products Affected

- Cometriq

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	

CABOZANTINIB S-MALATE - CABOMETYX

Products Affected

- Cabometyx oral tablet 20 mg, 40 mg, 60 mg

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PATIENT HAS RECEIVED PRIOR ANTIANGIOGENIC THERAPY (E.G., SUTENT [SUNITINIB], VOTRIENT [PAZOPANIB], INLYTA [AXITINIB], NEXAVAR [SORAFENIB])

CANAKINUMAB

Products Affected

- Ilaris (PF)

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	CAPS: 4 YEARS AND OLDER. SJIA: 2 YEARS AND OLDER.
Prescriber Restrictions	PRESCRIBED OR SUPERVISED BY RHEUMATOLOGIST, DERMATOLOGIST, OR AN IMMUNOLOGIST.
Coverage Duration	12 MONTHS
Other Criteria	

CANNABINOIDS

Products Affected

- dronabinol

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 MONTHS
Other Criteria	B VS D COVERAGE CONSIDERATION. PART D COVERAGE CONSIDERATION FOR A DIAGNOSIS OF NAUSEA AND VOMITING ASSOCIATED WITH CANCER CHEMOTHERAPY REQUIRES A TRIAL OF OR CONTRAINDICATION TO CONVENTIONAL ANTIEMETIC THERAPIES SUCH AS ONDANSETRON, STEROIDS INDICATED FOR EMESIS OR EMEND. NO ADDITIONAL REQUIREMENTS FOR A DIAGNOSIS OF ANOREXIA ASSOCIATED WITH WEIGHT LOSS IN PATIENTS WITH AIDS.

CANNABINOIDS ORAL SOLUTION

Products Affected

- Syndros

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 MONTHS
Other Criteria	B VS D COVERAGE CONSIDERATION. PART D COVERAGE CONSIDERATION FOR A DIAGNOSIS OF NAUSEA AND VOMITING ASSOCIATED WITH CANCER CHEMOTHERAPY REQUIRES A TRIAL OF OR CONTRAINDICATION TO FORMULARY DRONABINOL CAPSULES AND ONE CONVENTIONAL ANTIEMETIC THERAPY SUCH AS ONDANSETRON, STEROIDS INDICATED FOR EMESIS OR APREPITANT. PART D COVERAGE CONSIDERATION FOR A DIAGNOSIS OF ANOREXIA ASSOCIATED WITH WEIGHT LOSS IN PATIENTS WITH AIDS REQUIRES A TRIAL OF OR CONTRAINDICATION TO FORMULARY DRONABINOL CAPSULES.

CARFILZOMIB

Products Affected

- Kyprolis

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	

CERITINIB

Products Affected

- Zykadia

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	POSITIVE FOR ANAPLASTIC LYMPHOMA KINASE (ALK) FUSION ONCOGENE.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	

CERTOLIZUMAB PEGOL

Products Affected

- Cimzia
- Cimzia Powder for Reconst

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	RENEWAL: PHYSICIAN ATTESTATION OF IMPROVEMENT.
Age Restrictions	
Prescriber Restrictions	RHEUMATOID ARTHRITIS/ANKYLOSING SPONDYLITIS: PRESCRIBED BY OR GIVEN IN CONSULTATION WITH A RHEUMATOLOGIST. PSORIATIC ARTHRITIS: PRESCRIBED BY OR GIVEN IN CONSULTATION WITH A DERMATOLOGIST OR RHEUMATOLOGIST. CROHN'S DISEASE: PRESCRIBED BY OR GIVEN IN CONSULTATION WITH A GASTROENTEROLOGIST.
Coverage Duration	INITIAL: RA: 6 MONTHS. PSA/AS: 4 MONTHS. CD: 12 MONTHS. RENEWAL: 12 MONTHS FOR ALL DIAGNOSES.
Other Criteria	INITIAL: RHEUMATOID ARTHRITIS (RA): PREVIOUS TRIAL OF HUMIRA AND ONE DMARD (DISEASE-MODIFYING ANTIRHEUMATIC DRUG) SUCH AS METHOTREXATE, LEFLUNOMIDE, HYDROXYCHLOROQUINE, OR SULFASALAZINE. PSORIATIC ARTHRITIS (PSA): PREVIOUS TRIAL OF HUMIRA AND ONE DMARD (DISEASE-MODIFYING ANTIRHEUMATIC DRUG) SUCH AS METHOTREXATE, LEFLUNOMIDE, HYDROXYCHLOROQUINE, OR SULFASALAZINE. ANKYLOSING SPONDYLITIS: PREVIOUS TRIAL OF HUMIRA. CROHN'S DISEASE (CD): PREVIOUS TRIAL OF HUMIRA AND ONE CONVENTIONAL AGENT SUCH AS A CORTICOSTEROID (I.E., BUDESONIDE, METHYLPREDNISOLONE), AZATHIOPRINE, MERCAPTOPYRINE, METHOTREXATE, OR MESALAMINE.

CLOBAZAM

Products Affected

- Onfi oral suspension
- Onfi oral tablet 10 mg, 20 mg

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	2 YEARS OF AGE OR OLDER
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	TRIAL OF LAMOTRIGINE OR TOPIRAMATE. REQUESTS FOR ORAL SUSPENSION APPROVABLE IF PATIENT IS UNABLE TO SWALLOW OR IS UNDER THE AGE OF 5 YEARS.

COBIMETINIB FUMARATE

Products Affected

- Cotellic

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	

CORTICOTROPIN

Products Affected

- Acthar H.P.

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	INFANTILE SPASMS AND MULTIPLE SCLEROSIS: 28 DAYS. OTHER FDA APPROVED INDICATIONS:12 MONTHS.
Other Criteria	ALL FDA APPROVED INDICATIONS EXCEPT INFANTILE SPASMS REQUIRE A TRIAL OR CONTRAINDICATION TO IV CORTICOSTEROIDS. NOT APPROVED IN PATIENTS WITH ACUTE EXACERBATION OF MULTIPLE SCLEROSIS OR OTHER FDA APPROVED INDICATIONS IF IV ACCESS CAN BE OBTAINED. NOT APPROVED FOR DIAGNOSTIC PURPOSES.

CRIZOTINIB

Products Affected

- Xalkori

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	

DABRAFENIB MESYLATE

Products Affected

- Tafinlar

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	

DACLATASVIR

Products Affected

- Daklinza

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D. CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT AASLD-IDSA GUIDANCE AND ADDITIONAL CONSIDERATION FOR COVERAGE CONSISTENT WITH FDA LABELING.
Exclusion Criteria	
Required Medical Information	HCV RNA LEVEL WITHIN PAST 6 MONTHS.
Age Restrictions	
Prescriber Restrictions	GASTROENTEROLOGIST, INFECTIOUS DISEASE SPECIALIST, PHYSICIAN SPECIALIZING IN THE TREATMENT OF HEPATITIS (HEPATOLOGIST), OR A SPECIALLY TRAINED GROUP SUCH AS ECHO (EXTENSION FOR COMMUNITY HEALTHCARE OUTCOMES) MODEL.
Coverage Duration	CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT AASLD/IDSA GUIDANCE.

PA Criteria	Criteria Details
Other Criteria	<p>CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT AASLD/IDSA GUIDANCE. TRIAL OF A PREFERRED FORMULARY ALTERNATIVE INCLUDING HARVONI OR EPCLUSA WHEN THESE AGENTS ARE CONSIDERED ACCEPTABLE FOR TREATMENT OF THE SPECIFIC GENOTYPE PER AASLD/IDSA GUIDANCE. NO APPROVALS FOR CONCURRENT USE WITH ANY OF THESE (CONTRAINDICATED OR NOT RECOMMENDED BY THE MANUFACTURER) MEDICATIONS: AMIODARONE, CARBAMAZEPINE, PHENYTOIN, OR RIFAMPIN. APPROVAL FOR INTERFERON INELIGIBLE PATIENTS - INTERFERON INELIGIBILITY INCLUDES CONCURRENT DIAGNOSIS OF AUTOIMMUNE HEPATITIS OR OTHER AUTOIMMUNE DISORDER, A KNOWN HYPERSENSITIVITY REACTION (SUCH AS URTICARIA, ANGIOEDEMA, BRONCHOCONSTRICTION AND ANAPHYLAXIS TO ALPHA INTERFERONS, PEG, OR ANY COMPONENT OF PEGINTERFERON), DOCUMENTED DEPRESSION, DECOMPENSATED HEPATIC DISEASE: A BASELINE NEUTROPHIL COUNT BELOW 1,500 PER MICROLITER, A BASELINE PLATELET COUNT BELOW 90,000, OR A BASELINE HEMOGLOBIN BELOW 10G/DL THAT HAS NOT RESPONDED TO TREATMENT.</p>

DACLIZUMAB

Products Affected

- Zinbryta

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	PRE-EXISTING HEPATIC DISEASE OR IMPAIRMENT, INCLUDING: ACTIVE HEPATITIS B AND C, AUTOIMMUNE HEPATITIS OR OTHER AUTOIMMUNE CONDITIONS INVOLVING THE LIVER, BASELINE ALT AND AST GREATER THAN OR EQUAL TO 2 TIMES UPPER LIMIT OF NORMAL (ULN).
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	INITIAL: 3 MONTHS. RENEWAL: 12 MONTHS.
Other Criteria	PREVIOUS TRIAL OF TWO OF THE FOLLOWING PREFERRED AGENTS FOR MULTIPLE SCLEROSIS, SUCH AS AUBAGIO, AVONEX, GILENYA, PLEGRIDY, REBIF, TECFIDERA, OR FORMULARY GLATIRAMER ACETATE. RENEWAL: REQUESTS FOR DACLIZUMAB WILL NOT BE APPROVED FOR THE FOLLOWING PATIENTS: PATIENT WITH AUTOIMMUNE HEPATITIS OR HEPATIC INJURY.

DALFAMPRIDINE

Products Affected

- Ampyra

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	WALKING DISABILITY SUCH AS MILD TO MODERATE BILATERAL LOWER EXTREMITY WEAKNESS OR UNILATERAL WEAKNESS PLUS LOWER EXTREMITY OR TRUNCAL ATAXIA.
Age Restrictions	
Prescriber Restrictions	NEUROLOGIST
Coverage Duration	INITIAL: 3 MONTHS. RENEWAL: 12 MONTHS
Other Criteria	RENEWAL: PHYSICIAN ATTESTATION OF IMPROVEMENT IN WALKING ABILITY.

DARATUMUMAB

Products Affected

- Darzalex

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	

DASATINIB

Products Affected

- Sprycel oral tablet 100 mg, 140 mg, 20 mg, 50 mg, 70 mg, 80 mg

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PREVIOUSLY-TREATED CHRONIC MYELOID LEUKEMIA (CML) REQUIRES BCR-ABL MUTATIONAL ANALYSIS NEGATIVE FOR THE FOLLOWING MUTATIONS: T315I, V299L, T315A, F317L/V/I/C.

DEFERASIROX

Products Affected

- Exjade
- Jadenu
- Jadenu Sprinkle

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	PRESCRIBED BY OR GIVEN IN CONSULTATION WITH A HEMATOLOGIST OR HEMATOLOGIST/ONCOLOGIST
Coverage Duration	INITIAL: 6 MONTHS RENEWAL: 12 MONTHS
Other Criteria	CHRONIC IRON OVERLOAD DUE TO BLOOD TRANSFUSIONS INITIAL: SERUM FERRITIN LEVEL CONSISTENTLY ABOVE 1000 MCG/L. RENEWAL: SERUM FERRITIN LEVEL CONSISTENTLY ABOVE 500 MCG/L. NON-TRANSFUSION DEPENDENT THALASSEMIA (NTDT) INITIAL: SERUM FERRITIN LEVEL CONSISTENTLY ABOVE 300 MCG/L AND LIVER IRON CONCENTRATION (LIC) OF 5 MG FE/G DRY WEIGHT OR GREATER. RENEWAL: SERUM FERRITIN LEVEL CONSISTENTLY ABOVE THAN 300 MCG/L AND LIC OF 3 MG FE/G DRY WEIGHT OR GREATER

DEFERIPRONE

Products Affected

- Ferriprox

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	PRESCRIBED BY OR GIVEN IN CONSULTATION WITH A HEMATOLOGIST OR HEMATOLOGIST/ONCOLOGIST
Coverage Duration	INITIAL: 6 MONTHS RENEWAL: 12 MONTHS
Other Criteria	INITIAL CRITERIA: REQUIRES TRIAL OF EXJADE, JADENU, OR GENERIC DEFEROXAMINE AND ONE OF THE FOLLOWING CRITERIA 1) PHYSICIAN ATTESTATION THAT PATIENT IS EXPERIENCING INTOLERABLE TOXICITIES, CLINICALLY SIGNIFICANT ADVERSE EFFECTS, OR CONTRAINDICATION TO THESE THERAPIES OR 2) INADEQUATE CHELATION DEFINED BY ONE OF THE FOLLOWING: A) SERUM FERRITIN LEVEL CONSISTENTLY ABOVE 2500 MCG/L OR B) EVIDENCE OF CARDIAC IRON ACCUMULATION. RENEWAL: SERUM FERRITIN LEVELS MUST BE CONSISTENTLY ABOVE 500MCG/L

DEFEROXAMINE

Products Affected

- deferoxamine

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	AT LEAST 3 YEARS OF AGE OR OLDER
Prescriber Restrictions	PRESCRIBED BY OR GIVEN IN CONSULTATION WITH A HEMATOLOGIST OR HEMATOLOGIST/ONCOLOGIST
Coverage Duration	INITIAL: 6 MONTHS RENEWAL: 12 MONTHS
Other Criteria	INITIAL: SERUM FERRITIN LEVEL CONSISTENTLY ABOVE 1000MCG/L RENEWAL: SERUM FERRITIN LEVELS MUST BE CONSISTENTLY ABOVE 500MCG/L

DEFLAZACORT

Products Affected

- Emflaza oral suspension
- Emflaza oral tablet 18 mg, 30 mg, 36 mg, 6 mg

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	REPORTED INTOLERANCE TO PREDNISONE OR PREDNISOLONE IS A REPORTED INTOLERANCE IN THE EMFLAZA PRESCRIBING INFORMATION AS AN ADVERSE EVENT OF EMFLAZA
Required Medical Information	PHYSICIAN ATTESTATION OF GENETIC TESTING CONFIRMING DMD DIAGNOSIS.
Age Restrictions	
Prescriber Restrictions	PRESCRIBED BY OR GIVEN IN CONSULTATION WITH A NEUROLOGIST SPECIALIZING IN THE TREATMENT OF DMD.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS

PA Criteria	Criteria Details
Other Criteria	<p>INITIAL CRITERIA: REQUIRE TRIAL OF PREDNISON OR PREDNISOLONE FOR AT LEAST 6 MONTHS. REQUEST DUE TO LACK OF EFFICACY OF PREDNISON OR PREDNISOLONE AND ALL OF THE FOLLOWING CRITERIA ARE MET: 1)PATIENT IS NOT IN STAGE 1 (PRE-SYMPTOMATIC PHASE) 2) STEROID MYOPATHY HAS BEEN RULED OUT 3) PHYSICIAN ATTESTATION OF DETERIORATION IN AMBULATION, FUNCTIONAL STATUS, OR PULMONARY FUNCTION CONSISTENT WITH ADVANCING DISEASE USING STANDARD MEASURES [SUCH AS 6-MINUTE WALK DISTANCE (6MWD), TIME TO ASCEND/DESCEND 4 STAIRS, RISE FROM FLOOR TIME (GOWER'S MANEUVER), 10-METER RUN/WALK TIME, OR NORTH STAR AMBULATORY ASSESSMENT (NSAA), PHYSICIAN GLOBAL ASSESSMENTS (PGA), PULMONARY FUNCTION (FVC, PFTS), UPPER LIMB STRENGTH (PROPELLING A WHEELCHAIR 30 FEET)]. REQUEST DUE TO ADVERSE EFFECTS OF PREDNISON OR PREDNISOLONE THAT ARE NOT LISTED IN THE PRESCRIBING INFORMATION OF EMFLAZA REQUIRE PHYSICIAN ATTESTATION OF LITERATURE SUPPORTING EMFLAZA MITIGATES NAMED ADVERSE CONSEQUENCE.</p> <p>RENEWAL: APPROVAL FOR PATIENTS CURRENTLY AMBULATORY REQUIRES PHYSICIAN ATTESTATION OF FUNCTION, STABILIZATION, OR IMPROVEMENT IN STANDARD MEASURES SINCE TREATMENT WITH EMFLAZA THAT ARE BEING MONITORED, TRACKED, AND DOCUMENTED CONSISTENTLY. APPROVAL FOR PATIENTS CURRENTLY NON-AMBULATORY REQUIRES PHYSICIAN ATTESTATION OF MAINTENANCE OR LESS THAN EXPECTED DECLINE IN PULMONARY FUNCTION AND/OR UPPER LIMB STRENGTH ASSESSED BY STANDARD MEASURES (SUCH AS PULMONARY FUNCTION [FVC, PFTS], UPPER LIMB STRENGTH MEASURES [PROPELLING A WHEELCHAIR 30 FEET], PHYSICIAN GLOBAL ASSESSMENTS [PGA]) SINCE TREATMENT WITH EMFLAZA THAT ARE BEING MONITORED, TRACKED, AND DOCUMENTED CONSISTENTLY.</p>

DENOSUMAB-XGEVA

Products Affected

- Xgeva

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	DIAGNOSIS OF MULTIPLE MYELOMA
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	

DESIRUDIN

Products Affected

- Iprivask

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	1 MONTH
Other Criteria	

DEUTETRABENAZINE

Products Affected

- Austedo oral tablet 12 mg, 6 mg, 9 mg

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	

DICHLORPHENAMIDE

Products Affected

- Keveyis

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	HEPATIC INSUFFICIENCY, PULMONARY OBSTRUCTION, OR A HEALTH CONDITION THAT WARRANTS CONCURRENT USE OF HIGH-DOSE ASPIRIN
Required Medical Information	
Age Restrictions	18 YEARS AND OLDER
Prescriber Restrictions	
Coverage Duration	INITIAL: 2 MONTHS RENEWAL: 12 MONTHS
Other Criteria	RENEWAL REQUIRES THE PATIENT EXPERIENCED AT LEAST TWO FEWER ATTACKS PER WEEK FROM THEIR BASELINE

DICLOFENAC EPOLAMINE

Products Affected

- Flector

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION.

DICLOFENAC TOPICAL

Products Affected

- diclofenac sodium topical gel 3 %
- Pennsaid topical solution in metered-dose pump

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	SOLUTION AND GEL: 12 MONTHS
Other Criteria	

DIMETHYL FUMARATE

Products Affected

- Tecfidera oral capsule, delayed release(DR/EC) 120 mg, 120 mg (14)- 240 mg (46), 240 mg

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	

DINUTUXIMAB

Products Affected

- Unituxin

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	

DROXIDOPA

Products Affected

- Northera

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	BLOOD PRESSURE READINGS WHILE THE PATIENT IS SITTING AND ALSO WITHIN 3 MINUTES OF STANDING FROM A SUPINE (LYING FACE UP) POSITION AT BASELINE AND RENEWAL.
Age Restrictions	
Prescriber Restrictions	PRESCRIBED BY OR GIVEN IN CONSULTATION WITH A NEUROLOGIST OR CARDIOLOGIST.
Coverage Duration	INITIAL: 3 MONTHS RENEWAL: 12 MONTHS
Other Criteria	INITIAL: DIAGNOSIS OF ORTHOSTATIC HYPOTENSION AS DOCUMENTED BY A DECREASE OF AT LEAST 20 MMHG IN SYSTOLIC BLOOD PRESSURE OR 10 MMHG DIASTOLIC BLOOD PRESSURE WITHIN THREE MINUTES AFTER STANDING FROM A SITTING POSITION. RENEWAL: PATIENT HAD AN INCREASE IN SYSTOLIC BLOOD PRESSURE FROM BASELINE OF AT LEAST 10 MMHG UPON STANDING FROM A SUPINE (LYING FACE UP) POSITION.

DUPILUMAB

Products Affected

- Dupixent

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	INITIAL: PATIENT HAS MINIMUM BODY SURFACE AREA (BSA) INVOLVEMENT OF AT LEAST 10%, ECZEMA AREA AND SEVERITY INDEX (EASI) SCORE OF AT LEAST 16, OR PHYSICIAN GLOBAL ASSESSMENT (PGA) SCORE OF AT LEAST 3. RENEWAL: PHYSICIAN ATTESTATION OF IMPROVEMENT.
Age Restrictions	
Prescriber Restrictions	PRESCRIBED BY OR GIVEN IN CONSULTATION WITH A DERMATOLOGIST, ALLERGIST OR IMMUNOLOGIST
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL:12 MONTHS
Other Criteria	INITIAL: PREVIOUS TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING: TOPICAL CORTICOSTEROIDS, TOPICAL CALCINEURIN INHIBITORS [E.G., ELIDEL (PIMECROLIMUS), GENERIC TACROLIMUS OINTMENT], OR TOPICAL PDE4 INHIBITOR [E.G., EUCRISA (CRISABOROLE)].

DURVALUMAB

Products Affected

- Imfinzi

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	

EDARAVONE

Products Affected

- Radicava

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	

ELBASVIR/GRAZOPREVIR

Products Affected

- Zepatier

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D. CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT AASLD-IDSA GUIDANCE AND ADDITIONAL CONSIDERATION FOR COVERAGE CONSISTENT WITH FDA LABELING.
Exclusion Criteria	MODERATE OR SEVERE LIVER IMPAIRMENT (CHILD PUGH B OR C)
Required Medical Information	HCV RNA LEVEL WITHIN PAST 6 MONTHS. FOR GENOTYPE 1A -TESTING FOR NS5A RESISTANCE-ASSOCIATED POLYMORPHISMS.
Age Restrictions	
Prescriber Restrictions	GASTROENTEROLOGIST, INFECTIOUS DISEASE SPECIALIST, PHYSICIAN SPECIALIZING IN THE TREATMENT OF HEPATITIS (HEPATOLOGIST), OR A SPECIALLY TRAINED GROUP SUCH AS ECHO (EXTENSION FOR COMMUNITY HEALTHCARE OUTCOMES) MODEL.
Coverage Duration	CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT AASLD/IDSA GUIDANCE.

PA Criteria	Criteria Details
Other Criteria	<p>CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT AASLD/IDSA GUIDANCE. TRIAL OF A PREFERRED FORMULARY ALTERNATIVE INCLUDING HARVONI OR EPCLUSA WHEN THESE AGENTS ARE CONSIDERED ACCEPTABLE FOR TREATMENT OF THE SPECIFIC GENOTYPE PER AASLD/IDSA GUIDANCE. NO CONCURRENT USE OF SOVALDI AND ANY OF THE FOLLOWING AGENTS: PHENYTOIN, CARBAMAZEPINE, RIFAMPIN, EFAVIRENZ, ATAZANAVIR, DARUNAVIR, LOPINAVIR, SAQUINAVIR, TIPRANAVIR, CYCLOSPORINE, NAFCILLIN, KETOCONAZOLE, MODAFINIL, BOSENTAN, ETRAVIRINE, ELVITEGRAVIR/COBICISTAT/EMTRICITABINE/TENOFOVIR, ATORVASTATIN AT DOSES GREATER THAN 20MG PER DAY OR ROSUVASTATIN AT DOSES GREATER THAN 10MG PER DAY.</p>

ELIGLUSTAT TARTRATE

Products Affected

- Cerdelga

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	

ELOSULFASE ALFA

Products Affected

- Vimizim

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	LIFETIME OF MEMBERSHIP IN PLAN.
Other Criteria	

ELOTUZUMAB

Products Affected

- Empliciti

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	

ELTROMBOPAG

Products Affected

- Promacta

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	ITP:INITIAL: 2MO.RENEW:AFTER RESPONSE:12MO, INADEQUATE DOSE:2MO.HCV:12MO.SEVERE APLASTIC ANEMIA:12MO
Other Criteria	CHRONIC IMMUNE (IDIOPATHIC) THROMBOCYTOPENIA PURPURA (ITP): INITIAL: TRIAL OF OR CONTRAINDICATION TO CORTICOSTEROIDS, IMMUNOGLOBULINS, OR AN INSUFFICIENT RESPONSE TO SPLENECTOMY. ITP: RENEWAL: PATIENT HAS A CLINICAL RESPONSE AS DEFINED BY AN INCREASE IN PLATELET COUNT OF GREATER THAN OR EQUAL TO $50 \times 10^9/L$ (GREATER THAN OR EQUAL TO 50,000 PER UL) AT THE MAX DOSE OF 75MG PER DAY FOR 4 WEEKS. HEPATITIS C: CONCURRENT INTERFERON THERAPY.

ENDOTHELIN RECEPTOR ANTAGONISTS

Products Affected

- Letairis
- Opsumit
- Tracleer

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	DOCUMENTED CONFIRMATORY PULMONARY ARTERIAL HYPERTENSION (PAH) DIAGNOSIS BASED ON RIGHT HEART CATHETERIZATION. PATIENT HAS NYHA-WHO FUNCTIONAL CLASS II-IV SYMPTOMS.
Age Restrictions	
Prescriber Restrictions	PRESCRIBED BY OR IN CONSULTATION WITH A CARDIOLOGIST OR PULMONOLOGIST
Coverage Duration	INITIAL AND RENEWAL: 12 MONTHS
Other Criteria	INITIAL: MEAN PULMONARY ARTERY PRESSURE (PAP) OF AT LEAST 25 MMHG OR GREATER, PULMONARY CAPILLARY WEDGE PRESSURE (PCWP) OF 15 MMHG OR LESS, PULMONARY VASCULAR RESISTANCE (PVR) GREATER THAN 3 WOOD UNITS. LETAIRIS: PATIENT DOES NOT HAVE IDIOPATHIC PULMONARY FIBROSIS (IPF) TRACLEER: PATIENT DOES NOT HAVE ELEVATED LIVER ENZYMES (ALT, AST) MORE THAN 3 TIMES UPPER LIMIT OF NORMAL (ULN) OR INCREASES IN BILIRUBIN BY 2 OR MORE TIMES ULN. PATIENT IS NOT CONCURRENTLY TAKING CYCLOSPORINE A OR GLYBURIDE. RENEWAL: PATIENT SHOW IMPROVEMENT FROM BASELINE IN THE 6-MINUTE WALK DISTANCE OR PATIENT HAS A STABLE 6-MINUTE WALK DISTANCE WITH A STABLE/IMPROVED WHO FUNCTIONAL CLASS.

ENZALUTAMIDE

Products Affected

- Xtandi

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	TRIAL OF OR CONTRAINDICATION TO ZYTIGA (ABIRATERONE ACETATE) IS ALSO REQUIRED IN PATIENTS WHO DO NOT HAVE A CONTRAINDICATION OR INTOLERANCE TO PREDNISONE.

EPOPROSTENOL IV

Products Affected

- epoprostenol (glycine)

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	COVERED UNDER LOCAL COVERAGE POLICY OF APPLICABLE MEDICARE DMERC.
Required Medical Information	FORMULARY DRUG ADMINISTERED IN A LONG TERM CARE FACILITY TO A PATIENT WHOSE PART A COVERAGE HAS EXPIRED OR FORMULARY DRUG NOT ADMINISTERED VIA AN IMPLANTABLE PUMP OR AN EXTERNAL PUMP OR DRUG ADMINISTERED VIA AN IMPLANTABLE PUMP/AN EXTERNAL PUMP. DOCUMENTED CONFIRMATORY PULMONARY ARTERIAL HYPERTENSION (PAH) DIAGNOSIS BASED ON RIGHT HEART CATHETERIZATION. PATIENT HAS NYHA-WHO FUNCTIONAL CLASS III-IV SYMPTOMS.
Age Restrictions	
Prescriber Restrictions	PRESCRIBED BY OR GIVEN IN CONSULTATION WITH A CARDIOLOGIST OR PULMONOLOGIST.
Coverage Duration	INITIAL AND RENEWAL: 12 MONTHS
Other Criteria	INITIAL: MEAN PULMONARY ARTERY PRESSURE (PAP) OF AT LEAST 25 MMHG OR GREATER, PULMONARY CAPILLARY WEDGE PRESSURE (PCWP) OF 15 MMHG OR LESS, PULMONARY VASCULAR RESISTANCE (PVR) GREATER THAN 3 WOOD UNITS. RENEWAL: PATIENT HAS SHOWN IMPROVEMENT FROM BASELINE IN THE 6-MINUTE WALK DISTANCE OR PATIENT HAS A STABLE 6-MINUTE WALK DISTANCE WITH A STABLE/ IMPROVED WHO FUNCTIONAL CLASS.

ERLOTINIB

Products Affected

- Tarceva oral tablet 100 mg, 150 mg, 25 mg

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	

ERYTHROPOIESIS STIMULATING AGENTS - EPOETIN ALFA

Products Affected

- EPOGEN 10,000 UNITS/ML VIAL SDV, P/F, OUTER
- Epogen injection solution 2,000 unit/mL, 20,000 unit/2 mL, 20,000 unit/mL, 3,000 unit/mL, 4,000 unit/mL
- Procrit injection solution 10,000 unit/mL, 2,000 unit/mL, 20,000 unit/2 mL, 20,000 unit/mL, 3,000 unit/mL, 4,000 unit/mL, 40,000 unit/mL

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D. ADDITIONAL OFF LABEL ANEMIA IN HEPATITIS C BEING TREATED IN COMBINATION WITH RIBAVIRIN AND INTERFERON ALFA OR PEGINTERFERON ALFA.
Exclusion Criteria	

PA Criteria	Criteria Details
Required Medical Information	<p>INITIAL: CHRONIC RENAL FAILURE (CRF) AND ANEMIA RELATED TO ZIDOVUDINE THERAPY REQUIRES A HEMOGLOBIN LEVEL OF LESS THAN 10G/DL. CANCER CHEMOTHERAPY REQUIRES A HEMOGLOBIN LEVEL OF LESS THAN 10G/DL ANEMIA DUE TO CONCURRENT HEPATITIS C TREATMENT WITH RIBAVIRIN PLUS INTERFERON ALFA/PEGINTERFERON ALFA REQUIRES A HEMOGLOBIN LEVEL LESS THAN 10G/DL AND RIBAVIRIN DOSE REDUCTION (UNLESS CONTRAINDICATED).ELECTIVE NON-CARDIAC OR NON-VASCULAR SURGERY REQUIRES A HEMOGLOBIN LEVEL LESS THAN 13G/DL. RENEWAL: CHRONIC RENAL FAILURE REQUIRES THAT THE PATIENT MEETS ONE OF THE FOLLOWING: IF THE PATIENT IS CURRENTLY RECEIVING DIALYSIS TREATMENT: 1) HEMOGLOBIN LEVEL OF LESS THAN 11G/DL OR 2) HEMOGLOBIN LEVEL THAT HAS REACHED 11G/DL AND DOSE REDUCTION/INTERRUPTION IS REQUIRED TO REDUCE THE NEED FOR BLOOD TRANSFUSIONS. IF THE PATIENT IS NOT RECEIVING DIALYSIS TREATMENT: 1) HEMOGLOBIN LEVEL OF LESS THAN 10G/DL OR 2) HEMOGLOBIN LEVEL THAT HAS REACHED 10G/DL AND DOSE REDUCTION/INTERRUPTION IS REQUIRED TO REDUCE THE NEED FOR BLOOD TRANSFUSIONS. ANEMIA DUE TO CONCURRENT HEPATITIS C TREATMENT WITH RIBAVIRIN PLUS INTERFERON ALFA/PEGINTERFERON ALFA, OR ANEMIA DUE TO ZIDOVUDINE THERAPY REQUIRES HEMOGLOBIN LEVELS BETWEEN 10G/DL AND 12G/DL. ANEMIA DUE TO EFFECT OF CONCOMITANTLY ADMINISTERED CANCER CHEMOTHERAPY REQUIRES A HEMOGLOBIN LEVEL OF LESS THAN 10 G/DL OR THAT THE HEMOGLOBIN LEVEL DOES NOT EXCEED A LEVEL NEEDED TO AVOID RBC TRANSFUSION.</p>
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	<p>ANEMIA FROM MYELOSUPPRESSIVE CHEMO/CKD WITHOUT DIALYSIS/ZIDOVUDINE:12 MONTHS.SURGERY:1 MO.HCV:6 MOS.</p>

PA Criteria	Criteria Details
Other Criteria	ALL INDICATIONS: TRIAL OF PROCRIT. PART D MEMBER RECEIVING DIALYSIS OR IDENTIFIED AS A PART D END STAGE RENAL DISEASE MEMBER: PAYS UNDER PART B.

ERYTHROPOIESIS STIMULATING AGENTS - MIRCERA

Products Affected

- Mircera injection syringe 100 mcg/0.3 mL, 200 mcg/0.3 mL, 50 mcg/0.3 mL, 75 mcg/0.3 mL

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	INITIAL: CHRONIC RENAL FAILURE REQUIRES HEMOGLOBIN LEVELS LESS THAN 10G/DL RENEWAL: CHRONIC RENAL FAILURE REQUIRES THAT THE PATIENT MEETS ONE OF THE FOLLOWING: IF THE PATIENT IS CURRENTLY RECEIVING DIALYSIS TREATMENT: 1) HEMOGLOBIN LEVEL OF LESS THAN 11G/DL OR 2) HEMOGLOBIN LEVEL THAT HAS REACHED 11G/DL AND DOSE REDUCTION/INTERRUPTION IS REQUIRED TO REDUCE THE NEED FOR BLOOD TRANSFUSIONS. IF THE PATIENT IS NOT RECEIVING DIALYSIS TREATMENT: 1) HEMOGLOBIN LEVEL OF LESS THAN 10G/DL OR 2) HEMOGLOBIN LEVEL THAT HAS REACHED 10G/DL AND DOSE REDUCTION/INTERRUPTION IS REQUIRED TO REDUCE THE NEED FOR BLOOD TRANSFUSIONS.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	ANEMIA DUE TO CKD WITH OR WITHOUT DIALYSIS: 12 MONTHS.
Other Criteria	TRIAL OF PROCRIT. PART D MEMBER RECEIVING DIALYSIS OR IDENTIFIED AS A PART D END STAGE RENAL DISEASE MEMBER: PAYS UNDER PART B.

ETANERCEPT

Products Affected

- Enbrel
- Enbrel SureClick

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	INITIAL: PLAQUE PSORIASIS: MODERATE TO SEVERE PLAQUE PSORIASIS INVOLVING AT LEAST 5% BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, OR GENITAL AREA. RENEWAL: PHYSICIAN ATTESTATION OF IMPROVEMENT.
Age Restrictions	RHEUMATOID ARTHRITIS, ANKYLOSING SPONDYLITIS, PSORIATIC ARTHRITIS: 18 YEARS OR OLDER
Prescriber Restrictions	RHEUMATOID ARTHRITIS, POLYARTICULAR JUVENILE IDIOPATHIC ARTHRITIS, ANKYLOSING SPONDYLITIS: PRESCRIBED BY OR GIVEN IN CONSULTATION WITH A RHEUMATOLOGIST. PSORIATIC ARTHRITIS: PRESCRIBED BY OR GIVEN IN CONSULTATION WITH A DERMATOLOGIST OR RHEUMATOLOGIST. PLAQUE PSORIASIS: PRESCRIBED BY OR GIVEN IN CONSULTATION WITH A DERMATOLOGIST.
Coverage Duration	INITIAL: RA: 6 MONTHS. PJIA: 3 MONTHS. PSA/AS/PSO: 4 MONTHS. RENEWAL: 12 MONTHS FOR ALL DIAGNOSES

PA Criteria	Criteria Details
Other Criteria	<p>INITIAL: RHEUMATOID ARTHRITIS (RA): PREVIOUS TRIAL OF HUMIRA FOLLOWED BY ONE OF THE FOLLOWING PREFERRED AGENTS: ORENCIA, XELJANZ, CIMZIA, OR ACTEMRA. POLYARTICULAR JUVENILE IDIOPATHIC ARTHRITIS (PJIA): PREVIOUS TRIAL OF HUMIRA FOLLOWED BY ONE OF THE FOLLOWING PREFERRED AGENTS: ORENCIA OR ACTEMRA. PSORIATIC ARTHRITIS (PSA): PREVIOUS TRIAL OF HUMIRA FOLLOWED BY ONE OF THE FOLLOWING PREFERRED AGENTS: CIMZIA, OTEZLA, OR COSENTYX. ANKYLOSING SPONDYLITIS (AS): PREVIOUS TRIAL OF HUMIRA FOLLOWED BY ONE OF THE FOLLOWING PREFERRED AGENTS: CIMZIA OR COSENTYX. PLAQUE PSORIASIS (PSO): PREVIOUS TRIAL OF HUMIRA FOLLOWED BY ONE OF THE FOLLOWING PREFERRED AGENTS: COSENTYX OR OTEZLA.</p>

ETEPLIRSEN

Products Affected

- Exondys 51

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	PHYSICIAN ATTESTATION OF GENETIC TESTING CONFIRMING THAT MUTATION IN DUCHENNE MUSCULAR DYSTROPHY (DMD) GENE IS AMENABLE TO EXON 51 SKIPPING.
Age Restrictions	
Prescriber Restrictions	PRESCRIBED BY OR GIVEN IN CONSULTATION WITH A NEUROLOGIST SPECIALIZING IN THE TREATMENT OF DMD.
Coverage Duration	INITIAL: 24 WEEKS. RENEWAL: 12 MONTHS.
Other Criteria	INITIAL CRITERIA: PATIENT IS AMBULATORY AND IS CURRENTLY RECEIVING TREATMENT WITH OR HAS A CONTRAINDICATION TO CORTICOSTEROIDS. RENEWAL CRITERIA: PATIENT HAS MAINTAINED OR DEMONSTRATED A LESS THAN EXPECTED DECLINE IN AMBULATORY ABILITY IN MUSCLE FUNCTION ASSESSMENTS OR OTHER MUSCLE FUNCTION (I.E. PULMONARY OR CARDIAC FUNCTION) DURING THE PAST 24 WEEKS. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.

EVEROLIMUS

Products Affected

- Afinitor Disperz
- Afinitor oral tablet 10 mg, 2.5 mg, 5 mg, 7.5 mg

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	ADVANCED RENAL CELL CARCINOMA (RCC): TRIAL OF OR CONTRAINDICATION TO SUTENT OR NEXAVAR.

EVOLOCUMAB

Products Affected

- Repatha Pushtronex
- Repatha SureClick
- Repatha Syringe

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	HEFH OR ASCVD: 18 YEARS OF AGE AND OLDER.
Prescriber Restrictions	CARDIOLOGIST, ENDOCRINOLOGIST OR LIPIDOLOGIST
Coverage Duration	INITIAL: 6 MONTHS RENEWAL: 12 MONTHS

PA Criteria	Criteria Details
Other Criteria	<p>FOR HETEROZYGOUS FAMILIAL HYPERCHOLESTEROLEMIA (HEFH) OR ATHEROSCLEROTIC CARDIOVASCULAR DISEASE (ASCVD): MUST HAVE LDL LEVEL GREATER THAN 100MG/DL ON MAXIMAL DRUG TREATMENT (MDT) FOR AT LEAST 2 MONTHS WITHIN THE PAST 2 MONTHS AND ONE OF THE FOLLOWING: (1) HEFH DETERMINED BY SIMON BROOME DIAGNOSTIC (SBD) CRITERIA OR A SCORE OF 6 OR GREATER ON THE DUTCH LIPID NETWORK (DLN) CRITERIA OR (2) ASCVD AS SUBSTANTIATED BY PHYSICIAN ATTESTATION.</p> <p>HOMOZYGOUS FAMILIAL HYPERCHOLESTEROLEMIA (HOFH): LDL LEVEL GREATER THAN 100MG/DL ON MDT FOR AT LEAST 2 MONTHS WITHIN THE PAST 2 MONTHS AND HOFH DETERMINED BY ONE OF THE FOLLOWING: 1) SBD CRITERIA, 2) A SCORE OF 8 OR GREATER ON THE DLN CRITERIA, OR 3) A CLINICAL DIAGNOSIS BASED ON A HISTORY OF AN UNTREATED LDL-C CONCENTRATION GREATER THAN 500 MG/DL TOGETHER WITH EITHER XANTHOMA BEFORE 10 YEARS OF AGE, OR EVIDENCE OF HEFH IN BOTH PARENTS. NO CONCURRENT USE OF OTHER PCSK9 INHIBITORS. INITIAL THERAPY: FOR STATIN TOLERANT PATIENTS: MUST HAVE TRIED MAXIMALLY TOLERATED DOSE OF HIGH INTENSITY STATIN SUCH AS ATORVASTATIN OR ROSUVASTATIN. FOR STATIN INTOLERANT PATIENTS WITH HEFH OR ASCVD: ONE OF THE FOLLOWING MUST BE MET: PHYSICIAN ATTESTATION OF STATIN INTOLERANCE (INCLUDING BUT NOT LIMITED TO MYOPATHY), OR PATIENT HAS TRIED ROSUVASTATIN OR ATORVASTATIN AT ANY DOSE. PATIENTS WITH CONTRAINDICATIONS TO STATINS INCLUDING ACTIVE DECOMPENSATED LIVER DISEASE, NURSING FEMALE, PREGNANCY OR PLANS TO BECOME PREGNANT OR HYPERSENSITIVITY REACTIONS WILL BE APPROVED FOR REPATHA THERAPY WITHOUT DOCUMENTED STATIN INTOLERANCE. FOR STATIN INTOLERANT PATIENTS WITH HOFH: MUST BE ON MAX LIPID-LOWERING THERAPY INCLUDING ONE OF THE FOLLOWING: NIACIN, BILE ACID SEQUESTRANT, LOMITAPIDE OR MIPOMERSEN. QUALIFIERS MUST PROVIDE DOCUMENTATION OF STATIN INTOLERANCE TO ONE OF THE FOLLOWING: A HIGH INTENSITY STATIN (ROSUVASTATIN OR ATORVASTATIN) OR OTHER STATIN THERAPY AT ANY DOSE. STATIN INTOLERANT PATIENTS</p>

PA Criteria	Criteria Details
	MUST BE ON MAXIMAL LIPID-LOWERING MEDICATION (NON-STATIN THERAPY) FOR AT LEAST 2 MONTHS WITHIN THE PAST 2 MONTHS WITH DOCUMENTATION OF STATIN INTOLERANCE TO ATORVASTATIN OR ROSUVASTATIN OR STATIN THERAPY AT ANY DOSE. DOCUMENTATION OF STATIN INTOLERANCE INCLUDES: (1) PHYSICIAN ATTESTATION, OR (2) PATIENT HAS TRIED ROSUVASTATIN OR ATORVASTATIN AND HAS EXPERIENCED SKELETAL MUSCLE RELATED EVENTS (E.G. MYOPATHY). RENEWAL CRITERIA: RECEIVING PRIOR REPATHA THERAPY FOR AT LEAST 6 MONTHS AND NOT ON CONCURRENT THERAPY WITH OTHER PCSK9 INHIBITORS, MIPOMERSEN, OR LOMITAPIDE.

FENTANYL NASAL SPRAY

Products Affected

- Lazanda

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 MONTHS
Other Criteria	<p>CANCER: CURRENTLY ON A MAINTENANCE DOSE OF CONTROLLED-RELEASE OPIOID PAIN MEDICATION (SUCH AS MORPHINE SULFATE ER, OXYCODONE ER, OR FENTANYL). EITHER A TRIAL OR CONTRAINDICATION TO AT LEAST ONE (1) IMMEDIATE-RELEASE ORAL OPIOID PAIN AGENT (SUCH AS MORPHINE SULFATE IR, OXYCODONE/ASPIRIN, OXYCODONE/ACETAMINOPHEN, CODEINE/ACETAMINOPHEN, HYDROMORPHONE, OR MEPERIDINE) OR MEMBER HAS DIFFICULTY SWALLOWING TABLETS/CAPSULES AND TRIAL OR CONTRAINDICATION TO GENERIC FENTANYL CITRATE LOZENGE. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION.</p>

FENTANYL TRANSMUCOSAL AGENTS - FENTANYL CITRATE

Products Affected

- fentanyl citrate

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 MONTHS
Other Criteria	CANCER: CURRENTLY ON A MAINTENANCE DOSE OF CONTROLLED-RELEASE OPIOID PAIN MEDICATION (SUCH AS MORPHINE SULFATE ER, OXYCODONE ER, OR FENTANYL). EITHER A TRIAL OR CONTRAINDICATION TO AT LEAST ONE (1) IMMEDIATE-RELEASE ORAL OPIOID PAIN AGENT (SUCH AS MORPHINE SULFATE IR, OXYCODONE/ASPIRIN, OXYCODONE/ACETAMINOPHEN, CODEINE/ACETAMINOPHEN, HYDROMORPHONE, OR MEPERIDINE) OR MEMBER HAS DIFFICULTY SWALLOWING TABLETS/CAPSULES. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION.

FINGOLIMOD

Products Affected

- Gilenya

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	

GEFITINIB

Products Affected

- Iressa

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	

GLATIRAMER ACETATE

Products Affected

- Copaxone subcutaneous syringe 40 mg/mL
- Glatopa

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	

GLYCEROL PHENYL BUTYRATE

Products Affected

- Ravicti

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	TRIAL OF OR CONTRAINDICATION TO SODIUM PHENYL BUTYRATE (BUPHENYL).

GOLIMUMAB IV

Products Affected

- Simponi ARIA

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	RENEWAL: PHYSICIAN ATTESTATION OF IMPROVEMENT.
Age Restrictions	
Prescriber Restrictions	RHEUMATOID ARTHRITIS: PRESCRIBED BY OR GIVEN IN CONSULTATION WITH A RHEUMATOLOGIST
Coverage Duration	INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS
Other Criteria	INITIAL: PREVIOUS TRIAL OF HUMIRA AND ONE OF THE FOLLOWING PREFERRED AGENTS: ORENCIA, XELJANZ, CIMZIA, OR ACTEMRA.

GOLIMUMAB SQ

Products Affected

- Simponi

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	RENEWAL: PHYSICIAN ATTESTATION OF IMPROVEMENT.
Age Restrictions	
Prescriber Restrictions	RHEUMATOID ARTHRITIS, ANKYLOSING SPONDYLITIS: PRESCRIBED BY OR GIVEN IN CONSULTATION WITH A RHEUMATOLOGIST. PSORIATIC ARTHRITIS: PRESCRIBED BY OR GIVEN IN CONSULTATION WITH A DERMATOLOGIST OR RHEUMATOLOGIST. ULCERATIVE COLITIS: PRESCRIBED BY OR GIVEN IN CONSULTATION WITH A GASTROENTEROLOGIST.
Coverage Duration	INITIAL: RA: 6 MONTHS. PSA/AS: 4 MONTHS. UC: 12 MONTHS. RENEWAL: 12 MONTHS FOR ALL DIAGNOSES.
Other Criteria	INITIAL: RHEUMATOID ARTHRITIS (RA): PREVIOUS TRIAL OF HUMIRA AND ONE OF THE FOLLOWING PREFERRED AGENTS: ORENCIA, XELJANZ, CIMZIA OR ACTEMRA. PSORIATIC ARTHRITIS (PSA): PREVIOUS TRIAL OF HUMIRA AND ONE OF THE FOLLOWING PREFERRED AGENTS: CIMZIA, OTEZLA, OR COSENTYX. ANKYLOSING SPONDYLITIS (AS): PREVIOUS TRIAL OF HUMIRA AND ONE OF THE FOLLOWING PREFERRED AGENTS: CIMZIA OR COSENTYX. ULCERATIVE COLITIS (UC): PREVIOUS TRIAL OF HUMIRA AND ONE OF THE FOLLOWING CONVENTIONAL AGENTS SUCH AS A CORTICOSTEROID (I.E., BUDESONIDE, METHYLPREDINSONE), AZATHIOPRINE, MERCAPTOPYRINE, METHOTREXATE, OR MESALAMINE.

HIGH RISK DRUGS IN THE ELDERLY - ANTICHOLINERGICS - BENZTROPINE_TRIHEXYPHENIDYL

Products Affected

- benztropine
- trihexyphenidyl

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PRESCRIBER ACKNOWLEDGEMENT/AWARENESS DRUG IS LABELED AS HIGH RISK MEDICATION IN THE ELDERLY FOR PATIENTS 65 YEARS AND OLDER. HOSPICE PATIENTS WILL BE APPROVED WITHOUT REQUIRING PRESCRIBER ACKNOWLEDGEMENT.

HIGH RISK DRUGS IN THE ELDERLY - ANTICHOLINERGICS - PROMETHAZINE

Products Affected

- Phenadoz
- promethazine injection solution
- promethazine oral
- promethazine rectal
- Promethegan

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PRURITUS/URTICARIA/SEASONAL/PERENNIAL ALLERGY: TRIAL OR CONTRAINDICATION TO A NON-SEDATING ANTIHISTAMINE SUCH AS LEVOCETIRIZINE. NAUSEA AND VOMITING: PRESCRIBER ACKNOWLEDGEMENT OR AWARENESS THAT THE REQUESTED MEDICATION IS CONSIDERED HIGH-RISK FOR PATIENTS 65 YEARS AND OLDER. HOSPICE PATIENTS WILL BE APPROVED WITHOUT TRIAL OF FORMULARY ALTERNATIVES NOR REQUIRING PRESCRIBER ACKNOWLEDGEMENT.

HIGH RISK DRUGS IN THE ELDERLY - ANTI- INFECTIVE

Products Affected

- nitrofurantoin
- nitrofurantoin macrocrystal
- nitrofurantoin monohyd/m-cryst

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS. PA REQUIRED FOR PATIENTS 65 YEARS AND OLDER WITH OVER 90 DAYS CUMULATIVE USE OF THE REQUESTED AGENT.
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PREVIOUS TRIAL OF OR CONTRAINDICATION TO SULFAMETHOXAZOLE/TRIMETHOPRIM (TMP-SMX) OR TRIMETHOPRIM. HOSPICE PATIENTS ARE APPROVED WITHOUT REQUIRING A PREVIOUS TRIAL OF FORMULARY ALTERNATIVES.

HIGH RISK DRUGS IN THE ELDERLY - BARBITURATE COMBINATIONS

Products Affected

- Ascomp with Codeine
- Butalbital Compound W/Codeine
- butalbital-acetaminop-caff-cod
- butalbital-acetaminophen oral tablet 50-325 mg
- butalbital-acetaminophen-caff oral capsule 50-325-40 mg
- butalbital-acetaminophen-caff oral tablet 50-325-40 mg
- butalbital-aspirin-caffeine oral capsule
- Capacet
- Tencon oral tablet 50-325 mg
- Zebutal oral capsule 50-325-40 mg

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PRESCRIBER ACKNOWLEDGEMENT/AWARENESS DRUG IS LABELED AS HIGH RISK MEDICATION IN THE ELDERLY FOR PATIENTS 65 YEARS AND OLDER. HOSPICE PATIENTS ARE APPROVED WITHOUT REQUIRING PRESCRIBER ACKNOWLEDGEMENT.

HIGH RISK DRUGS IN THE ELDERLY - CARDIOVASCULAR

Products Affected

- guanfacine oral tablet
- nifedipine oral capsule

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	HYPERTENSION: PREVIOUS TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING GENERIC FORMULARY ALTERNATIVES: ANGIOTENSIN CONVERTING ENZYME INHIBITOR (ACE INHIBITOR), ACE INHIBITOR COMBINATION, ANGIOTENSIN RECEPTOR BLOCKER (ARB), ARB COMBINATION, BETA BLOCKER, BETA BLOCKER COMBINATION, OR CALCIUM CHANNEL BLOCKERS. PRESCRIBER ACKNOWLEDGEMENT/AWARENESS THAT THE DRUG IS CONSIDERED HIGH RISK FOR PATIENTS 65 YEARS AND OLDER. HOSPICE PATIENTS WILL BE APPROVED WITHOUT REQUIRING A PREVIOUS TRIAL OF FORMULARY ALTERNATIVES OR PRESCRIBER ACKNOWLEDGEMENT.

HIGH RISK DRUGS IN THE ELDERLY - DIGOXIN

Products Affected

- Digitek oral tablet 125 mcg, 250 mcg
- DIGOXIN 0.25 MG/ML SYRINGE
- digoxin injection solution
- digoxin oral solution 50 mcg/mL
- digoxin oral tablet 125 mcg, 250 mcg
- Lanoxin oral tablet 187.5 mcg, 62.5 mcg

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	DIGOXIN LEVEL
Age Restrictions	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	APPROVAL FOR MEMBERS STABLE ON DOSES GREATER THAN 125 MCG PER DAY WITH DOCUMENTED THERAPEUTIC DIGOXIN LEVEL TAKEN WITHIN THE PAST YEAR. HOSPICE PATIENTS ARE APPROVED WITHOUT REQUIRING DIGOXIN LEVELS.

HIGH RISK DRUGS IN THE ELDERLY - ENDOCRINE - ESTROGEN

Products Affected

- CombiPatch
- Duavee
- estradiol oral
- estradiol transdermal patch semiweekly
- estradiol transdermal patch weekly
- estradiol-norethindrone acet
- estropipate
- Fyavolv oral tablet 1-5 mg-mcg
- Jinteli
- Lopreeza
- Menest oral tablet 0.3 mg, 0.625 mg, 1.25 mg
- Mimvey
- Mimvey Lo
- norethindrone ac-eth estradiol oral tablet 0.5-2.5 mg-mcg, 1-5 mg-mcg
- Premarin oral
- Premphase
- Prempro

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
Prescriber Restrictions	
Coverage Duration	12 MONTHS

PA Criteria	Criteria Details
Other Criteria	VULVAR/VAGINAL ATROPHY, OSTEOPOROSIS AND VASOMOTOR SYMPTOMS OF MENOPAUSE: PRESCRIBER ACKNOWLEDGEMENT/AWARENESS THAT THE DRUG IS CONSIDERED HIGH RISK FOR PATIENTS 65 YEARS AND OLDER. ALL OTHER FDA APPROVED INDICATIONS NOT PREVIOUSLY MENTIONED IN THIS SECTION, SUCH AS PALLIATIVE TREATMENT, AND HOSPICE PATIENTS ARE APPROVED WITHOUT REQUIRING PRESCRIBER ACKNOWLEDGEMENT.

HIGH RISK DRUGS IN THE ELDERLY - ENDOCRINE - SULFONYLUREAS

Products Affected

- glyburide
- glyburide-metformin
- glyburide micronized

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	TRIAL OF GLIMEPIRIDE, GLIPIZIDE, OR PRESCRIBER ACKNOWLEDGEMENT/AWARENESS THAT THE DRUG IS CONSIDERED HIGH RISK FOR PATIENTS 65 YEARS AND OLDER. HOSPICE PATIENTS ARE APPROVED WITHOUT A TRIAL OF FORMULARY ALTERNATIVES OR PRESCRIBER ACKNOWLEDGEMENT.

HIGH RISK DRUGS IN THE ELDERLY - NON-BENZODIAZEPINE

Products Affected

- eszopiclone
- zolpidem oral
- zaleplon

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS. PA REQUIRED FOR PATIENTS 65 YEARS AND OLDER WITH OVER 90 DAYS CUMULATIVE USE OF NON-BENZODIAZEPINE AGENTS.
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	TRIAL OF SILENOR AND BELSOMRA OR PRESCRIBER ACKNOWLEDGEMENT/ AWARENESS THAT THE DRUG IS CONSIDERED HIGH RISK FOR PATIENTS 65 YEARS AND OLDER. HOSPICE PATIENTS ARE APPROVED WITHOUT REQUIRING A TRIAL OF FORMULARY ALTERNATIVES (SILENOR AND BELSOMRA) OR PRESCRIBER ACKNOWLEDGEMENT.

HIGH RISK DRUGS IN THE ELDERLY - SKELETAL MUSCLE RELAXANTS

Products Affected

- carisoprodol
- chlorzoxazone
- cyclobenzaprine oral tablet
- methocarbamol oral

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PRESCRIBER ACKNOWLEDGEMENT/AWARENESS DRUG IS LABELED AS HIGH RISK MEDICATION IN THE ELDERLY FOR PATIENTS 65 YEARS AND OLDER. HOSPICE PATIENTS ARE APPROVED WITHOUT REQUIRING PRESCRIBER ACKNOWLEDGEMENT.

HIGH RISK DRUGS IN THE ELDERLY- ANTICHOLINERGICS- CYPROHEPTADINE

Products Affected

- cyproheptadine

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PRESCRIBER ACKNOWLEDGEMENT/AWARENESS DRUG IS CONSIDERED A HIGH RISK MEDICATION IN THE ELDERLY FOR PATIENTS 65 YEARS AND OLDER. HOSPICE PATIENTS ARE APPROVED WITHOUT REQUIRING PRESCRIBER ACKNOWLEDGEMENT.

HIGH RISK DRUGS IN THE ELDERLY- ANTICHOLINERGICS- DIPHENHYDRAMINE ELIXIR

Products Affected

- diphenhydramine HCl oral elixir

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	ANTIHISTAMINIC CONDITIONS (PRURITUS OR URTICARIA): TRIAL OR CONTRAINDICATION TO A NON-SEDATING ANTIHISTAMINE SUCH AS LEVOCETIRIZINE. INSOMNIA: TRIAL OF SILENOR AND BELSOMRA. ANTIPARKINSONISM: PRESCRIBER ACKNOWLEDGEMENT/AWARENESS DRUG IS CONSIDERED AS HIGH RISK MEDICATION IN THE ELDERLY FOR PATIENTS 65 YEARS AND OLDER. HOSPICE PATIENTS WILL BE APPROVED WITHOUT TRIAL OF FORMULARY ALTERNATIVES NOR REQUIRING PRESCRIBER ACKNOWLEDGEMENT.

HIGH RISK DRUGS IN THE ELDERLY- DIPHENOXYLATE-ATROPINE

Products Affected

- diphenoxyate-atropine

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PRESCRIBER ACKNOWLEDGEMENT/AWARENESS THAT THE DRUG IS CONSIDERED HIGH RISK FOR PATIENTS 65 YEARS AND OLDER. HOSPICE PATIENTS WILL BE APPROVED WITHOUT REQUIRING PRESCRIBER ACKNOWLEDGEMENT.

HIGH RISK DRUGS IN THE ELDERLY- HYDROXYZINE

Products Affected

- hydroxyzine HCl intramuscular
- hydroxyzine HCl oral solution 10 mg/5 mL
- hydroxyzine HCl oral tablet
- hydroxyzine pamoate

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PRESCRIBER ACKNOWLEDGEMENT/AWARENESS THAT THE DRUG IS CONSIDERED HIGH RISK FOR PATIENTS 65 YEARS AND OLDER. HOSPICE PATIENTS REQUIRE PHYSICIAN ATTESTATION THAT REQUESTED MEDICATION IS USED TO TREAT A DIAGNOSIS UNRELATED TO THE TERMINAL ILLNESS OR RELATED CONDITION, AND ARE APPROVED WITHOUT REQUIRING PRESCRIBER ACKNOWLEDGEMENT.

HIGH RISK DRUGS IN THE ELDERLY- INDOMETHACIN

Products Affected

- indomethacin oral capsule 25 mg, 50 mg
- indomethacin oral capsule, extended release

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PREScriBER ACKNOWLEDGEMENT/AWARENESS DRUG IS LABELED AS HIGH RISK MEDICATION IN THE ELDERLY FOR PATIENTS 65 YEARS AND OLDER. HOSPICE PATIENTS WILL BE APPROVED WITHOUT REQUIRING PREScriBER ACKNOWLEDGEMENT.

HIGH RISK DRUGS IN THE ELDERLY- KETOROLAC ORAL

Products Affected

- ketorolac oral

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
Prescriber Restrictions	
Coverage Duration	30 DAYS
Other Criteria	PRESCRIBER ACKNOWLEDGEMENT/AWARENESS THAT THE DRUG IS CONSIDERED HIGH RISK FOR PATIENTS 65 YEARS AND OLDER. HOSPICE PATIENTS WILL BE APPROVED WITHOUT REQUIRING PRESCRIBER ACKNOWLEDGEMENT.

HIGH RISK DRUGS IN THE ELDERLY- MECLIZINE

Products Affected

- meclizine oral tablet 12.5 mg, 25 mg

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PREScriBER ACKNOWLEDGEMENT/AWARENESS DRUG IS CONSIDERED HIGH RISK MEDICATION IN THE ELDERLY FOR PATIENTS 65 YEARS AND OLDER. FOR NAUSEA, VOMITING, AND DIZZINESS ASSOCIATED WITH MOTION SICKNESS: TRIAL OF OR CONTRAINDICATION TO PROCHLORPERAZINE, PROCHLORPERAZINE MALEATE, OR PROCHLORPERAZINE EDISYLATE. HOSPICE PATIENTS REQUIRE PHYSICIAN ATTESTATION THAT REQUESTED MEDICATION IS USED TO TREAT A DIAGNOSIS UNRELATED TO THE TERMINAL ILLNESS OR RELATED CONDITION, AND ARE APPROVED WITHOUT TRIAL OF FORMULARY ALTERNATIVE OR REQUIRING PREScriBER ACKNOWLEDGEMENT.

HIGH RISK DRUGS IN THE ELDERLY- MEGESTROL

Products Affected

- megestrol oral suspension 400 mg/10 mL
- megestrol oral tablet (40 mg/mL)

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PRESCRIBER ACKNOWLEDGEMENT/AWARENESS THAT THE DRUG IS CONSIDERED HIGH RISK FOR PATIENTS 65 YEARS AND OLDER. HOSPICE PATIENTS WILL BE APPROVED WITHOUT REQUIRING PRESCRIBER ACKNOWLEDGEMENT.

HIGH RISK DRUGS IN THE ELDERLY- PAROXETINE

Products Affected

- paroxetine HCl oral tablet
- paroxetine HCl oral tablet extended release 24 hr
- Paxil oral suspension

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PRESCRIBER ACKNOWLEDGEMENT/AWARENESS THAT THE DRUG IS CONSIDERED HIGH RISK FOR PATIENTS 65 YEARS AND OLDER. HOSPICE PATIENTS WILL BE APPROVED WITHOUT REQUIRING PRESCRIBER ACKNOWLEDGEMENT.

HIGH RISK DRUGS IN THE ELDERLY- TCA

Products Affected

- amitriptyline
- amitriptyline-chlordiazepoxide
- amoxapine
- clomipramine
- desipramine
- doxepin oral
- imipramine HCl
- imipramine pamoate
- nortriptyline
- perphenazine-amitriptyline
- protriptyline
- Surmontil
- trimipramine

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PRESCRIBER ACKNOWLEDGEMENT/AWARENESS THAT THE DRUG IS CONSIDERED HIGH RISK FOR PATIENTS 65 YEARS AND OLDER. HOSPICE PATIENTS WILL BE APPROVED WITHOUT REQUIRING PRESCRIBER ACKNOWLEDGEMENT. PRIOR AUTHORIZATION APPLIES TO NEW START ONLY.

HIGH RISK DRUGS IN THE ELDERLY- BENZODIAZEPINE SEDATIVE HYPNOTICS

Products Affected

- estazolam oral tablet 1 mg, 2 mg
- flurazepam oral capsule 15 mg, 30 mg
- temazepam oral capsule 15 mg, 30 mg
- triazolam oral tablet 0.125 mg, 0.25 mg

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS. PA REQUIRED FOR PATIENTS 65 YEARS AND OLDER WITH OVER 90 DAYS CUMULATIVE USE OF THE REQUESTED AGENT.
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PREVIOUS TRIAL OF OR CONTRAINDICATION TO SILENOR AND BELSOMRA OR PRESCRIBER ACKNOWLEDGEMENT/AWARENESS THAT THE DRUG IS LABELED AS HIGH RISK FOR PATIENTS 65 YEARS AND OLDER. HOSPICE PATIENTS WILL BE APPROVED WITHOUT REQUIRING A TRIAL OF FORMULARY ALTERNATIVES OR PRESCRIBER ACKNOWLEDGEMENT.

HIGH RISK MEDICATIONS IN THE ELDERLY- PHENOBARBITAL

Products Affected

- phenobarbital

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PRESCRIBER ACKNOWLEDGEMENT/AWARENESS THAT THE DRUG IS CONSIDERED HIGH RISK FOR PATIENTS 65 YEARS AND OLDER. FOR TREATMENT OF EPILEPSY/SEIZURES IN PATIENTS WHO ARE NOT CURRENTLY STABLE ON PHENOBARBITAL: PATIENT HAS NOT RESPONDED TO OTHER ANTICONVULSANTS. FOR SHORT TERM INSOMNIA: PREVIOUS TRIAL OF OR CONTRAINDICATION TO BELSOMRA AND SILENOR. PATIENTS WHO ARE STABLE ON PHENOBARBITAL FOR EPILEPSY/SEIZURES OR HOSPICE PATIENTS ARE APPROVED WITHOUT REQUIRING A TRIAL OF FORMULARY ALTERNATIVES OR PRESCRIBER ACKNOWLEDGEMENT.

HYDROXYPROGESTERONE CAPROATE- DELALUTIN GENERIC

Products Affected

- hydroxyprogesterone caproate

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.

IBRUTINIB

Products Affected

- Imbruvica

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	

IBUPROFEN-FAMOTIDINE

Products Affected

- Duexis

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	TRIAL OF ONE OF THE FOLLOWING GENERIC, FEDERAL LEGEND HISTAMINE H2-RECEPTOR ANTAGONISTS: FAMOTIDINE, CIMETIDINE, NIZATIDINE, OR RANITIDINE, AND TRIAL OF GENERIC, FEDERAL LEGEND IBUPROFEN.

IDELALISIB

Products Affected

- Zydelig

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	

IMATINIB MESYLATE

Products Affected

- imatinib oral tablet 100 mg, 400 mg

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	ALL DIAGNOSES: 12 MONTHS. ADJUVANT GASTROINTESTINAL STROMAL TUMOR (GIST) TREATMENT: 36 MONTHS.
Other Criteria	PATIENTS WITH PREVIOUSLY-TREATED CML REQUIRE A BCR-ABL MUTATIONAL ANALYSIS CONFIRMING THAT THE PATIENT IS NEGATIVE FOR THE FOLLOWING MUTATIONS: T315I, V299L, F317L/V/I/C, Y253H, E255K/V, F359V/C/I.

IMIQUIMOD - ALDARA

Products Affected

- imiquimod

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	ACTINIC KERATOSIS: PRESCRIBED BY OR GIVEN IN CONSULTATION WITH A DERMATOLOGIST. SUPERFICIAL BASAL CELL CARCINOMA: PRESCRIBED BY OR GIVEN IN CONSULTATION WITH A DERMATOLOGIST OR AN ONCOLOGIST.
Coverage Duration	4 MONTHS
Other Criteria	EXTERNAL GENITAL WARTS: TRIAL OF PODOFILOX (CONDYLOX) 0.5% TOPICAL SOLUTION. ACTINIC KERATOSIS BRAND DRUG REQUEST: TRIAL OF GENERIC IMIQUIMOD 5% CREAM. SUPERFICIAL BASAL CELL CARCINOMA: LESS THAN 2CM IN SIZE AND NOT ON THE FACE.

INFLIXIMAB

Products Affected

- Remicade

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	INITIAL: PLAQUE PSORIASIS: MODERATE TO SEVERE PLAQUE PSORIASIS INVOLVING GREATER THAN OR EQUAL TO 5 % BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, OR GENITAL AREA. RENEWAL: PHYSICIAN ATTESTATION OF IMPROVEMENT.
Age Restrictions	
Prescriber Restrictions	RHEUMATOID ARTHRITIS, ANKYLOSING SPONDYLITIS: PRESCRIBED BY OR GIVEN IN CONSULTATION WITH A RHEUMATOLOGIST. PSORIATIC ARTHRITIS PRESCRIBED BY OR GIVEN IN CONSULTATION WITH A DERMATOLOGIST OR RHEUMATOLOGIST. PSORIASIS PRESCRIBED BY OR GIVEN IN CONSULTATION WITH A DERMATOLOGIST. CROHN'S DISEASE/ULCERATIVE COLITIS: PRESCRIBED BY OR GIVEN IN CONSULTATION WITH A GASTROENTEROLOGIST.
Coverage Duration	INITIAL: CD/UC: 8 MO. RA: 6 MO. PSA/AS/PSO: 4 MO. RENEWAL FOR ALL INDICATIONS: 12 MO.

PA Criteria	Criteria Details
Other Criteria	<p>INITIAL: RHEUMATOID ARTHRITIS (RA): PREVIOUS TRIAL OF HUMIRA FOLLOWED BY ONE OF THE FOLLOWING: ORENCIA, XELJANZ, CIMZIA, OR ACTEMRA. PSORIATRIC ARTHRITIS (PSA): PREVIOUS TRIAL OF HUMIRA FOLLOWED BY ONE OF THE FOLLOWING: CIMZIA, OTEZLA, OR COSENTYX. ANKYLOSING SPONDYLITIS (AS): PREVIOUS TRIAL OF HUMIRA FOLLOWED BY CIMZIA OR COSENTYX. PLAQUE PSORIASIS (PSO): PREVIOUS TRIAL OF HUMIRA FOLLOWED BY ONE OF THE FOLLOWING: COSENTYX OR OTEZLA. CROHN'S DISEASE (CD): PREVIOUS TRIAL OF HUMIRA FOLLOWED BY CIMZIA. ULCERATIVE COLITIS (UC): PREVIOUS TRIAL OF HUMIRA FOLLOWED BY ONE OF THE FOLLOWING: SIMPONI OR ENTYVIO.</p>

INFLIXIMAB-DYYB

Products Affected

- Inflectra

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	INITIAL: PLAQUE PSORIASIS: SEVERE PLAQUE PSORIASIS INVOLVING GREATER THAN OR EQUAL TO 5 PERCENT BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, OR GENITAL AREA. RENEWAL: PHYSICIAN ATTESTATION OF IMPROVEMENT.
Age Restrictions	
Prescriber Restrictions	RHEUMATOID ARTHRITIS, ANKYLOSING SPONDYLITIS: PRESCRIBED BY GIVEN OR IN CONSULTATION WITH A RHEUMATOLOGIST. PSORIATIC ARTHRITIS: PRESCRIBED BY OR GIVEN IN CONSULTATION WITH A DERMATOLOGIST OR RHEUMATOLOGIST. PSORIASIS: PRESCRIBED BY OR GIVEN IN CONSULTATION WITH A DERMATOLOGIST. CROHN'S DISEASE/ULCERATIVE COLITIS: GASTROENTEROLOGIST.
Coverage Duration	INITIAL: CD/UC: 8 MOS. RA: 6 MOS. PSA/AS/PSO: 4 MOS. RENEWAL FOR ALL INDICATIONS: 12 MOS.

PA Criteria	Criteria Details
Other Criteria	<p>INITIAL: RHEUMATOID ARTHRITIS (RA): PREVIOUS TRIAL OF HUMIRA FOLLOWED BY ONE OF THE FOLLOWING: ORENCIA, XELJANZ, CIMZIA, OR ACTEMRA. PSORIATIC ARTHRITIS (PSA): PREVIOUS TRIAL OF HUMIRA FOLLOWED BY ONE OF THE FOLLOWING: CIMZIA, OTEZLA, OR COSENTYX. ANKYLOSING SPONDYLITIS (AS): PREVIOUS TRIAL OF HUMIRA FOLLOWED BY ONE OF THE FOLLOWING: CIMZIA OR COSENTYX. PLAQUE PSORIASIS (PSO): PREVIOUS TRIAL OF HUMIRA FOLLOWED BY ONE OF THE FOLLOWING: COSENTYX OR OTEZLA. CROHN'S DISEASE (CD): PREVIOUS TRIAL OF HUMIRA FOLLOWED BY CIMZIA. ULCERATIVE COLITIS (UC): PREVIOUS TRIAL OF HUMIRA FOLLOWED BY ONE OF THE FOLLOWING: SIMPONI OR ENTYVIO.</p>

INTERFERON ALFA-2B

Products Affected

- Intron A injection

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	HEPATITIS C: GASTROENTEROLOGIST, INFECTIOUS DISEASE SPECIALIST, PHYSICIAN SPECIALIZING IN THE TREATMENT OF HEPATITIS (E.G. HEPATOLOGIST). NO REQUIREMENT FOR OTHER FDA APPROVED INDICATIONS.
Coverage Duration	6 MONTHS
Other Criteria	LIMITED TO 1 YEAR OF THERAPY EXCEPT 18 MONTHS FOR FOLLICULAR LYMPHOMA. HEPATITIS C GENOTYPE 1, 2, 3, 4, 5, OR 6: REQUIRES A TRIAL OF OR CONTRAINDICATION TO PEGINTERFERON ALFA-2A OR PEGINTERFERON ALFA-2B USED IN COMBINATION WITH RIBAVIRIN UNLESS CONTRAINDICATED.

INTERFERONS FOR MS-AVONEX, PLEGRIDY, REBIF

Products Affected

- Avonex (with albumin)
- Avonex intramuscular pen injector kit
- Avonex intramuscular syringe kit
- Plegridy
- Rebif (with albumin)
- Rebif Rebidose
- Rebif Titration Pack

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	

INTERFERONS FOR MS-BETASERON, EXTAVIA

Products Affected

- Betaseron subcutaneous kit
- Extavia subcutaneous kit

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	TRIAL WITH TWO OF THE FOLLOWING AGENTS FOR MULTIPLE SCLEROSIS: AUBAGIO, AVONEX, GILENYA, PLEGRIDY, REBIF, TECFIDERA, AND GLATIRAMER
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	

IPILIMUMAB

Products Affected

- Yervoy

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	INITIAL: UNRESECTABLE/METASTATIC MELANOMA: 3 MO ADJVNT MELANOMA: 6 MO RENEWAL: ADJVNT MELANOMA: 6 MO
Other Criteria	RENEWAL FOR ADJUVANT CUTANEOUS MELANOMA: NO EVIDENCE OF DISEASE RECURRENCE (DEFINED AS THE APPEARANCE OF ONE OR MORE NEW MELANOMA LESIONS: LOCAL, REGIONAL OR DISTANT METASTASIS).

IVABRADINE

Products Affected

- Corlanor

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	PATIENT MUST HAVE NEW YORK HEART ASSOCIATION (NYHA) CLASS II TO IV HEART FAILURE
Age Restrictions	
Prescriber Restrictions	PRESCRIBED BY OR GIVEN IN CONSULTATION WITH A CARDIOLOGIST.
Coverage Duration	INITIAL AND RENEWAL: 12 MONTHS.
Other Criteria	INITIAL: APPROVAL REQUIRES THE PATIENT DOES NOT HAVE A DEMAND PACEMAKER SET TO A RATE OF 60 BEATS PER MINUTE OR GREATER. PATIENT IS CURRENTLY RECEIVING TREATMENT WITH OR HAS AN INTOLERANCE TO A FORMULARY BETA BLOCKER SUCH AS METOPROLOL SUCCINATE, BISOPROLOL, OR CARVEDILOL. RENEWAL: APPROVAL REQUIRES DIAGNOSIS OF HEART FAILURE AND PATIENT MUST BE IN SINUS RHYTHM.

IVACAFTOR

Products Affected

- Kalydeco

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	HOMOZYGOUS FOR F508DEL MUTATION IN CFTR GENE.
Required Medical Information	CONFIRMED MUTATION IN CFTR GENE ACCEPTABLE FOR THE TREATMENT OF CYSTIC FIBROSIS.
Age Restrictions	6 YEARS OF AGE OR OLDER.
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	

IVACAFTOR - GRANULE PACKETS

Products Affected

- Kalydeco

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	F508DEL MUTATION IN CFTR GENE.
Required Medical Information	CONFIRMED MUTATION IN CFTR GENE ACCEPTABLE FOR THE TREATMENT OF CYSTIC FIBROSIS. PATIENT WEIGHT.
Age Restrictions	2 YEARS OF AGE TO 5 YEARS OF AGE
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	

IXAZOMIB

Products Affected

- Ninlaro

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	

IXEKIZUMAB

Products Affected

- Taltz Autoinjector
- Taltz Syringe

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	PLAQUE PSORIASIS (PSO): MODERATE TO SEVERE PLAQUE PSORIASIS INVOLVING GREATER THAN OR EQUAL TO 5PERCENT BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, OR GENITAL AREA. RENEWAL: PHYSICIAN ATTESTATION OF IMPROVEMENT.
Age Restrictions	
Prescriber Restrictions	PLAQUE PSORIASIS (PSO): PRESCRIBED BY OR GIVEN IN CONSULTATION WITH A DERMATOLOGIST
Coverage Duration	PLAQUE PSORIASIS (PSO): INITIAL: 4 MONTHS. RENEWAL: 12 MONTHS
Other Criteria	INITIAL: PLAQUE PSORIASIS (PSO): PREVIOUS TRIAL OF HUMIRA FOLLOWED BY ONE OF THE FOLLOWING PREFERRED AGENTS: COSENTYX OR OTEZLA.

LEDIPASVIR-SOFOSBUVIR

Products Affected

- Harvoni

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D. CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT AASLD-IDSA GUIDANCE AND ADDITIONAL CONSIDERATION FOR COVERAGE CONSISTENT WITH FDA LABELING.
Exclusion Criteria	
Required Medical Information	HCV RNA LEVEL WITHIN PAST 6 MONTHS.
Age Restrictions	
Prescriber Restrictions	GASTROENTEROLOGIST, INFECTIOUS DISEASE SPECIALIST, PHYSICIAN SPECIALIZING IN THE TREATMENT OF HEPATITIS (HEPATOLOGIST), OR A SPECIALLY TRAINED GROUP SUCH AS ECHO (EXTENSION FOR COMMUNITY HEALTHCARE OUTCOMES) MODEL.
Coverage Duration	CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT AASLD/IDSA GUIDANCE.
Other Criteria	CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT AASLD/IDSA GUIDANCE. PATIENT IS NOT CONCURRENTLY TAKING ANY OF THE FOLLOWING: CARBAMAZEPINE, PHENYTOIN, PHENOBARBITAL, OXCARBAZEPINE, RIFAMPIN, RIFABUTIN, RIFAPENTINE, ROSUVASTATIN, SIMEPREVIR, SOFOSBUVIR (AS A SINGLE AGENT), STRIBILD (ELVITEGRAVIR/COBICISTAT/EMTRICITABINE /TENOFIVIR), OR TIPRANA VIR/RITONAVIR.

LENALIDOMIDE

Products Affected

- Revlimid

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	

LENVATINIB MESYLATE

Products Affected

- Lenvima

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	

LIDOCAINE

Products Affected

- lidocaine topical adhesive patch,medicated
- lidocaine topical ointment

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D. ADDITIONAL COVERAGE FOR DIABETIC NEUROPATHY WILL BE CONSIDERED FOR REQUESTS FOR LIDOCAINE TOPICAL PATCHES.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	PATCH: 12 MONTHS. OINTMENT: 3 MONTHS.
Other Criteria	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION.

LIFITEGRAST OPHTHALMIC

Products Affected

- Xiidra

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	PRESCRIBED BY OR GIVEN IN CONSULTATION WITH AN OPHTHALMOLOGIST, OPTOMETRIST OR RHEUMATOLOGIST.
Coverage Duration	12 MONTHS
Other Criteria	PREVIOUS TRIAL OF OR CONTRAINDICATION TO CYCLOSPORINE OPHTHALMIC (RESTASIS).

LOMITAPIDE

Products Affected

- Juxtapid oral capsule 10 mg, 20 mg, 30 mg, 40 mg, 5 mg, 60 mg

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	LDL CHOLESTEROL LEVEL, LDL RECEPTOR STATUS
Age Restrictions	
Prescriber Restrictions	PRESCRIBED BY OR IN CONSULTATION WITH A CARDIOLOGIST, ENDOCRINOLOGIST OR LIPIDOLOGIST.
Coverage Duration	INITIAL: 7 MONTHS RENEWAL: 6 MONTHS

PA Criteria	Criteria Details
Other Criteria	<p>DIAGNOSIS CONFIRMED BY ONE OF THE FOLLOWING: SIMON BROOME DIAGNOSTIC CRITERIA (DEFINITE), (E.G. GENETIC TESTING CONSISTENT WITH HOFH AND PRETREATMENT BASELINE LDL CHOLESTEROL IS GREATER THAN 190 MG/DL), CASCADE SCREENING, DUTCH LIPID NETWORK CRITERIA WITH A SCORE OF AT LEAST 6, OR HISTORY OF UNTREATED CHOLESTEROL GREATER THAN 500MG/DL (OR TREATED CHOLESTEROL GREATER THAN 300MG/DL) AND CUTANEOUS XANTHOMA BEFORE 10 YEARS OF AGE. LOMITAPIDE WILL NOT BE APPROVED FOR PATIENTS CONCURRENTLY USING ANY OF THE FOLLOWING STRONG OR MODERATE CYP3A4 MEDICATIONS: CLARITHROMYCIN, CONIVAPTAN, INDINAVIR, ITRACONAZOLE, KETOCONAZOLE, LOPINAVIR/RITONAVIR, MIBEFRADIL, NEFAZODONE, NELFINAVIR, POSACONAZOLE, RITONAVIR, SAQUINAVIR, TELITHROMYCIN, TIPRANAVIR/RITONAVIR, VORICONAZOLE, AMPRENAVIR, APREPITANT, ATAZANAVIR, CIPROFLOXACIN, CRIZOTINIB, DARUNAVIR/RITONAVIR, DILTIAZEM, ERYTHROMYCIN, FLUCONAZOLE, FOSAMPRENAVIR, IMATINIB, OR VERAPAMIL. INITIAL: LDL CHOLESTEROL LEVEL OF AT LEAST 160MG/DL WHILE ON LIPID-LOWERING THERAPY PRIOR TO INITIATING LOMITAPIDE. PREVIOUS TRIAL OF A PCSK9 INHIBITOR (E.G. ALIROCUMAB OR EVOLOCUMAB), UNLESS THE PATIENT HAS NON-FUNCTIONING LDL RECEPTORS. PREVIOUS TRIAL OF ROSUVASTATIN OR ATORVASTATIN, UNLESS THE PATIENT HAS AN ABSOLUTE CONTRAINDICATION TO STATIN THERAPY (E.G. ACTIVE, DECOMPENSATED LIVER DISEASE, NURSING FEMALE, PREGNANCY OR PLANS TO BECOME PREGNANT, HYPERSENSITIVITY REACTION). STATIN-TOLERANT PATIENTS MUST BE TAKING ATORVASTATIN OR ROSUVASTATIN FOR THE PAST 2 MONTHS PRIOR TO STARTING LOMITAPIDE. LOMITAPIDE MUST BE USED IN COMBINATION WITH ATORVASTATIN OR ROSUVASTATIN. IF THE PATIENT HAS PREVIOUSLY TRIED ATORVASTATIN OR ROSUVASTATIN, LOMITAPIDE MUST BE USED IN COMBINATION WITH ANOTHER STATIN OR FORMULARY LDL-LOWERING AGENT (E.G. BILE ACID SEQUESTRANT, GEMFIBROZIL OR OTHER FIBRATE, EZETIMIBE, OR NIACIN). STATIN-INTOLERANT PATIENTS REQUIRE EITHER PHYSICIAN ATTESTATION OF</p>

PA Criteria	Criteria Details
	<p>STATIN INTOLERANCE OR HISTORY OF SKELETAL-MUSCLE RELATED SYMPTOMS (E.G., MYOPATHY) DUE TO A PREVIOUS TRIAL OF STATINS (E.G. ROSUVASTATIN OR ATORVASTATIN). FOR STATIN-INTOLERANT PATIENTS, LOMITAPIDE MUST BE USED IN COMBINATION WITH ONE OF THE FOLLOWING FORMULARY LIPID-LOWERING TREATMENTS: EZETIMIBE, FENOFIBRATE, NIACIN, OR A BILE ACID SEQUESTRANT (E.G. CHOLESTYRAMINE, COLESTIPOL, COLESEVELAM). RENEWAL: PATIENT HAS RECEIVED AT LEAST 6 MONTHS OF THERAPY WITH LOMITAPIDE IN COMBINATION WITH ANOTHER AND LIPID-LOWERING AGENT.</p>

LUMACAFITOR-IVACAFITOR

Products Affected

- Orkambi

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	CONFIRMED MUTATION IN CFTR GENE ACCEPTABLE FOR THE TREATMENT OF CYSTIC FIBROSIS. BASELINE FEV1.
Age Restrictions	
Prescriber Restrictions	PRESCRIBED BY OR GIVEN IN CONSULTATION WITH A PULMONOLOGIST OR CYSTIC FIBROSIS EXPERT.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL 12 MONTHS.
Other Criteria	RENEWAL: MAINTAINED OR IMPROVEMENT IN FEV1 OR REDUCTION IN NUMBER OF PULMONARY EXACERBATIONS OR IMPROVEMENT IN BODY MASS INDEX (BMI).

MEPOLIZUMAB

Products Affected

- Nucala

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	CONCURRENT USE OF XOLAIR
Required Medical Information	BLOOD EOSINOPHIL LEVEL GREATER THAN OR EQUAL TO 150 CELLS/MCL WITHIN THE LAST 6 WEEKS OR GREATER THAN OR EQUAL TO 300 CELLS/MCL WITHIN THE LAST 12 MONTHS
Age Restrictions	
Prescriber Restrictions	PRESCRIBED BY OR IN CONSULTATION WITH A PHYSICIAN SPECIALIZING IN PULMONARY MEDICINE, AN ALLERGIST OR AN IMMUNOLOGIST.
Coverage Duration	INITIAL 24 WEEKS. RENEWAL 12 MONTHS
Other Criteria	INITIAL THERAPY: PATIENT CURRENTLY TREATED WITH A MAXIMALLY TOLERATED DOSE OF INHALED CORTICOSTEROIDS AND AT LEAST ONE OTHER MAINTENANCE MEDICATION WHICH INCLUDES ANY OF THE FOLLOWING: LONG-ACTING INHALED BETA2-AGONIST, LONG-ACTING MUSCARINIC ANTAGONIST, A LEUKOTRIENE RECEPTOR ANTAGONIST, THEOPHYLLINE, OR ORAL CORTICOSTEROID. RENEWAL REQUIRES DOCUMENTATION THAT THE PATIENT HAS EXPERIENCED IMPROVEMENT IN ASTHMA EXACERBATIONS FROM BASELINE (PHYSICIAN ATTESTATION) AND A REDUCTION IN ORAL CORTICOSTEROID DOSE (IF THE PATIENT WAS ON A MAINTENANCE REGIMEN OF ORAL CORTICOSTEROIDS AT THE INITIATION OF TREATMENT).

METHYLNALTREXONE

Products Affected

- Relistor subcutaneous solution
- Relistor subcutaneous syringe

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	ADVANCED ILLNESS: OPIOID-INDUCED CONSTIPATION.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 MONTHS FOR PATIENTS RECEIVING PALLIATIVE CARE, 12 MONTHS FOR PATIENTS WITH CHRONIC, NON-CANCER PAIN.
Other Criteria	ADVANCED ILLNESS: PATIENT IS RECEIVING PALLIATIVE CARE. CHRONIC NON-CANCER PAIN: PATIENT HAS BEEN TAKING OPIOIDS FOR AT LEAST 4 WEEKS AND HAD A PREVIOUS TRIAL OF OR CONTRAINDICATION TO NALOXEGOL (MOVANTIK).

METHYLNALTREXONE ORAL

Products Affected

- Relistor oral

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PATIENT HAS BEEN TAKING OPIOIDS FOR AT LEAST 4 WEEKS AND HAD A PREVIOUS TRIAL OF OR CONTRAINDICATION TO NALOXEGOL (MOVANTIK).

MIDOSTAURIN

Products Affected

- Rydapt

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	ACUTE MYELOID LEUKEMIA: 6 MONTHS. ADVANCED SYSTEMIC MASTOCYTOSIS: 12 MONTHS
Other Criteria	

MIFEPRISTONE

Products Affected

- Korlym

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	

MILTEFOSINE

Products Affected

- Impavido

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	

MIPOMERSEN

Products Affected

- Kynamro

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	LDL CHOLESTEROL LEVEL, LDL RECEPTOR STATUS
Age Restrictions	
Prescriber Restrictions	PRESCRIBED BY OR GIVEN IN CONSULTATION WITH A CARDIOLOGIST, ENDOCRINOLOGIST OR LIPIDOLOGIST
Coverage Duration	INITIAL: 7 MONTHS RENEWAL 12 MONTHS

PA Criteria	Criteria Details
Other Criteria	<p>DIAGNOSIS OF HOMOZYGOUS FAMILIAL HYPERCHOLESTEROLEMIA AS DETERMINED BY ONE OF THE FOLLOWING CRITERIA: SIMON BROOME DIAGNOSTIC CRITERIA (DEFINITE) [EXAMPLE: GENETIC TESTING CONSISTENT WITH HOFH AND PRETREATMENT BASELINE LDL CHOLESTEROL IS GREATER THAN 190 MG/DL], CASCADE SCREENING, DUTCH LIPID NETWORK CRITERIA WITH A SCORE AT LEAST 6, OR HISTORY OF UNTREATED CHOLESTEROL GREATER THAN 500MG/DL (OR TREATED GREATER THAN 300MG/DL) AND CUTANEOUS XANTHOMA BEFORE AGE 10. INITIAL CRITERIA: CURRENT LDL CHOLESTEROL LEVEL IS AT LEAST 160MG/DL. PATIENT DOES NOT HAVE ANY OF THE FOLLOWING CONTRAINDICATIONS TO KYNAMRO (MIPOMERSEN): MODERATE OR SEVERE HEPATIC IMPAIRMENT OR ACTIVE LIVER DISEASE, INCLUDING UNEXPLAINED PERSISTENT ELEVATIONS OF SERUM TRANSAMINASES. PREVIOUS TRIAL OF A PCSK9 INHIBITOR (SUCH AS ALIROCUMAB OR EVOLOCUMAB) UNLESS THE PATIENT HAS NON-FUNCTIONING LDL RECEPTORS. PREVIOUS TRIAL WITH ONE OF THE FOLLOWING STATINS: ROSUVASTATIN OR ATORVASTATIN. PATIENTS WITH ABSOLUTE CONTRAINDICATION TO STATIN THERAPY (ACTIVE, DECOMPENSATED LIVER DISEASE, NURSING FEMALE, PREGNANCY OR PLANS TO BECOME PREGNANT, HYPERSENSITIVITY REACTION) WILL BE APPROVED FOR THERAPY WITHOUT REQUIREMENT OF A TRIAL WITH A STATIN. STATIN-TOLERANT PATIENTS: PRIOR TO (KYNAMRO), PATIENT MUST HAVE BEEN TAKING ONE OF THE FOLLOWING: ATORVASTATIN OR ROSUVASTATIN, FOR AT LEAST 2 MONTHS WITHIN THE PAST 2 MONTHS. FOR STATIN-INTOLERANT PATIENTS: DOCUMENTATION OF STATIN INTOLERANCE WHICH INCLUDES THE FOLLOWING: PHYSICIAN ATTESTATION OR PATIENT HAS TRIED ROSUVASTATIN OR ATORVASTATIN AND HAS EXPERIENCED SKELETAL-MUSCLE RELATED SYMPTOMS (E.G., MYOPATHY). UNLESS CONTRAINDICATED, PATIENT MUST BE ON CONCURRENT THERAPY WITH ONE OF THE FOLLOWING LIPID-LOWERING TREATMENTS (SUCH AS A STATIN [SIMVASTATIN, ATORVASTATIN], EZETIMIBE, FENOFIBRATE, NIACIN, OR BILE ACID SEQUESTRANT [CHOLESTYRAMINE, COLESTIPOL, COLESEVELAM]). RENEWAL CRITERIA:</p>

PA Criteria	Criteria Details
	PATIENT HAS RECEIVED THERAPY FOR AT LEAST 6 MONTHS AND MUST ALSO BE TAKING KYNAMRO IN COMBINATION WITH ANOTHER LIPID-LOWERING AGENT.

MODAFINIL AND ARMODAFINIL

Products Affected

- armodafinil

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION.

NAPROXEN- ESOMEPRAZOLE

Products Affected

- Vimovo

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	TRIAL OF ONE OF THE FOLLOWING GENERIC, FEDERAL LEGEND PROTON PUMP INHIBITORS: OMEPRAZOLE, LANSOPRAZOLE, OR PANTOPRAZOLE AND A TRIAL OF GENERIC, FEDERAL LEGEND NAPROXEN.

NATALIZUMAB

Products Affected

- Tysabri

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	CROHN'S DISEASE: PRESCRIBED BY OR GIVEN IN CONSULTATION WITH A GASTROENTEROLOGIST.
Coverage Duration	MULTIPLE SCLEROSIS: 12 MONTHS. CROHN'S DISEASE: INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS
Other Criteria	MULTIPLE SCLEROSIS INITIAL CRITERIA: PREVIOUS TRIAL OF TWO OF THE FOLLOWING PREFERRED AGENTS FOR MULTIPLE SCLEROSIS: GLATIRAMER, REBIF, AVONEX, PLEGRIDY, TECFIDERA, GILENYA, OR AUBAGIO. CROHN'S DISEASE INITIAL CRITERIA: PREVIOUS TRIAL OF HUMIRA AND CIMZIA. CROHN'S DISEASE RENEWAL CRITERIA: PATIENT HAS RECEIVED AT LEAST 12 MONTHS OF THERAPY WITH TYSABRI WITH PHYSICIAN ATTESTATION THAT THE PATIENT HAS NOT REQUIRED MORE THAN 3 MONTHS OF CORTICOSTEROID USE WITHIN THE PAST 12 MONTHS TO CONTROL THEIR CROHN'S DISEASE WHILE ON TYSABRI, OR PATIENT HAS ONLY RECEIVED 6 MONTHS OF THERAPY WITH TYSABRI WITH PHYSICIAN ATTESTATION THAT THE PATIENT HAS TAPERED OFF CORTICOSTEROIDS DURING THE FIRST 24 WEEKS OF TYSABRI THERAPY.

NECITUMUMAB

Products Affected

- Portrazza

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	

NILOTINIB

Products Affected

- Tasigna

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PREVIOUSLY TREATED CML REQUIRES BCR-ABL MUTATIONAL ANALYSIS NEGATIVE FOR THE FOLLOWING MUTATIONS: T315I, Y253H, E255K/V, AND F359V/C/I.

NINTEDANIB

Products Affected

- Ofev

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	NOT APPROVED FOR PATIENTS WITH OTHER KNOWN CAUSES OF INTERSTITIAL LUNG DISEASE (E.G., CONNECTIVE TISSUE DISEASE, DRUG TOXICITY, ASBESTOS OR BERYLLIUM EXPOSURE, HYPERSENSITIVITY PNEUMONITIS, SYSTEMIC SCLEROSIS, RHEUMATOID ARTHRITIS, RADIATION, SARCOIDOSIS, BRONCHIOLITIS OBLITERANS ORGANIZING PNEUMONIA, HUMAN IMMUNODEFICIENCY VIRUS (HIV) INFECTION, VIRAL HEPATITIS, AND CANCER. NOT APPROVED IF PATIENT DOES NOT HAVE A PREDICTED FORCED VITAL CAPACITY (FVC) OF AT LEAST 50 PERCENT OR HAS NOT OBTAINED LIVER FUNCTION TESTS
Required Medical Information	A USUAL INTERSTITIAL PNEUMONIA (UIP) PATTERN AS EVIDENCED BY HIGH-RESOLUTION COMPUTED TOMOGRAPHY (HRCT) ALONE OR VIA A COMBINATION OF SURGICAL LUNG BIOPSY AND HRCT.
Age Restrictions	
Prescriber Restrictions	PRESCRIBED BY OR GIVEN IN CONSULTATION WITH A PULMONOLOGIST
Coverage Duration	12 MONTHS
Other Criteria	

NIRAPARIB TOSYLATE

Products Affected

- Zejula

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	

NITISINONE

Products Affected

- Orfadin

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	DIAGNOSIS OF HEREDITARY TYROSINEMIA TYPE 1 AS CONFIRMED BY ELEVATED URINARY OR PLASMA SUCCINYLACETONE LEVELS OR A MUTATION IN THE FUMARYLACETOACETATE HYDROLASE GENE.
Age Restrictions	
Prescriber Restrictions	PRESCRIBED BY OR GIVEN IN CONSULTATION WITH A PRESCRIBER SPECIALIZING IN INHERITED METABOLIC DISEASES.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS
Other Criteria	INITIAL: PATIENT MUST BE COUNSELED ON MAINTAINING DIETARY RESTRICTION OF TYROSINE AND PHENYLALANINE (CONSISTENT WITH FDA LABELING). ORFADIN SUSPENSION: TRIAL OF ORFADIN CAPSULES. RENEWAL: THE PATIENT'S URINARY OR PLASMA SUCCINYLACETONE LEVELS HAVE DECREASED FROM BASELINE WHILE ON TREATMENT WITH NITISINONE.

NIVOLUMAB

Products Affected

- Opdivo

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	MELANOMA: OPDIVO IS NOT APPROVED FOR COMBINATION THERAPY WITH TAFINLAR, MEKINIST (TRAMETINIB), COTELLIC (COBIMETINIB), OR ZELBORAF.

OBETICHOLIC ACID

Products Affected

- Ocaliva

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	PATIENTS WITH COMPLETE BILIARY OBSTRUCTION.
Required Medical Information	DIAGNOSIS OF PRIMARY BILIARY CHOLANGITIS AS CONFIRMED BY AT LEAST TWO OF THE FOLLOWING CRITERIA: AN ALKALINE PHOSPHATASE LEVEL OF AT LEAST 1.5 TIMES THE UPPER LIMIT OF NORMAL (ULN), THE PRESENCE OF ANTIMITOCHONDRIAL ANTIBODIES AT A TITER OF 1:40 OR HIGHER, HISTOLOGIC EVIDENCE OF NON-SUPPURATIVE DESTRUCTIVE CHOLANGITIS AND DESTRUCTION OF INTERLOBULAR BILE DUCTS.
Age Restrictions	
Prescriber Restrictions	PRESCRIBED BY OR GIVEN IN CONSULTATION WITH A GASTROENTEROLOGIST OR HEPATOLOGIST.
Coverage Duration	12 MONTHS
Other Criteria	INITIAL: USED IN COMBINATION WITH URSODEOXYCHOLIC ACID (E.G., URSODIOL, URSO 250, URSO FORTE) IN ADULTS WITH AN INADEQUATE RESPONSE TO URSODEOXYCHOLIC ACID AT A DOSAGE OF 13-15 MG/KG/DAY FOR AT LEAST 1 YEAR, OR AS MONOTHERAPY IN ADULTS UNABLE TO TOLERATE URSODEOXYCHOLIC ACID. RENEWAL: PATIENT'S ALKALINE PHOSPHATASE LEVELS HAVE DECREASED BY AT LEAST 15% FROM BASELINE WHILE ON TREATMENT WITH OBETICHOLIC ACID.

OBINUTUZUMAB

Products Affected

- Gazyva

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 MONTHS
Other Criteria	

OCRELIZUMAB

Products Affected

- Ocrevus

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	RELAPSING FORM OF MULTIPLE SCLEROSIS: TRIAL OF TWO OF THE FOLLOWING AGENTS FOR MULTIPLE SCLEROSIS: AUBAGIO, AVONEX, GILENYA, PLEGRIDY, REBIF, TECFIDERA, OR GLATIRAMER. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.

OLAPARIB

Products Affected

- Lynparza

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	

OLARATUMAB

Products Affected

- Lartruvo

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	THIS DRUG MAY BE COVERED UNDER MEDICARE PART B OR D DEPENDING UPON THE CIRCUMSTANCES. INFORMATION MAY NEED TO BE SUBMITTED DESCRIBING THE USE AND SETTING OF THE DRUG TO MAKE THE DETERMINATION.

OMACETAXINE

Products Affected

- Synribo

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	INDUCTION: 3 MONTHS. POST INDUCTION OR RENEWAL: 3 TO 12 MONTHS
Other Criteria	CML INDUCTION THERAPY: TRIAL OF OR CONTRAINDICATION TO AT LEAST TWO OF THE FOLLOWING AGENTS: GLEEVEC, SPRYCEL, TASIGNA, BOSULIF OR ICLUSIG. APPROVAL FOR POST-INDUCTION THERAPY DURATION WILL DEPEND ON THE PATIENT'S HEMATOLOGIC RESPONSE, DEFINED AS AN ABSOLUTE NEUTROPHIL COUNT (ANC) GREATER THAN OR EQUAL TO $1.5 \times 10^9/L$, PLATELETS GREATER THAN OR EQUAL TO $100 \times 10^9/L$ WITHOUT BLOOD BLASTS OR THE PATIENT HAS BONE MARROW BLASTS AT LESS THAN 5 PERCENT. APPROVAL IS FOR 12 MONTHS IF HEMATOLOGIC RESPONSE IS MET. IF NOT MET, APPROVAL IS FOR 3 MONTHS.

OMALIZUMAB

Products Affected

- Xolair

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	INITIAL CRITERIA FOR ASTHMA: PATIENT MEETS THE CRITERIA OF MODERATE TO SEVERE ASTHMA, POSITIVE SKIN PRICK OR RAST TEST, FEV1 LESS THAN 80%, DEMONSTRATED INADEQUATELY CONTROLLED SYMPTOMS ON INHALED CORTICOSTEROIDS AND SECOND ASTHMA CONTROLLER, BASELINE IGE SERUM LEVEL GREATER THAN OR EQUAL TO 30IU/ML. RENEWAL CRITERIA FOR ASTHMA: PHYSICIAN ATTESTATION OF IMPROVEMENT IN ASTHMA EXCERBATIONS FROM BASELINE AND A REDUCTION IN ORAL OR INHALED CORTICOSTEROID USE.
Age Restrictions	
Prescriber Restrictions	PRESCRIBED BY OR GIVEN IN CONSULTATION WITH A SPECIALIST IN ALLERGY, PULMONARY MEDICINE, DERMATOLOGY OR IMMUNOLOGY.
Coverage Duration	ASTHMA: 12 MONTHS. CHRONIC IDIOPATHIC URTICARIA: 6 MONTHS.
Other Criteria	FOR CHRONIC IDIOPATHIC URTICARIA: PREVIOUS TRIAL OF OR CONTRAINDICATION TO A HIGH DOSE H1 ANTI-HISTAMINE (SUCH AS CLARINEX OR XYZAL) FOR AT LEAST 2 WEEKS AND STILL EXPERIENCE HIVES ON MOST DAYS OF THE WEEK.

OMBITASVIR-PARITAPREVIR-RITONAVIR

Products Affected

- Technivie

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D. CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT AASLD-IDSA GUIDANCE AND ADDITIONAL CONSIDERATION FOR COVERAGE CONSISTENT WITH FDA LABELING.
Exclusion Criteria	CIRRHOSIS, MODERATE OR SEVERE LIVER IMPAIRMENT (CHILD-PUGH B OR C).
Required Medical Information	HCV RNA LEVEL WITHIN PAST 6 MONTHS.
Age Restrictions	
Prescriber Restrictions	GASTROENTEROLOGIST, INFECTIOUS DISEASE SPECIALIST, PHYSICIAN SPECIALIZING IN THE TREATMENT OF HEPATITIS (HEPATOLOGIST), OR A SPECIALLY TRAINED GROUP SUCH AS ECHO (EXTENSION FOR COMMUNITY HEALTHCARE OUTCOMES) MODEL.
Coverage Duration	CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT AASLD/IDSA GUIDANCE.

PA Criteria	Criteria Details
Other Criteria	<p>CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT AASLD/IDSA GUIDANCE. TRIAL OF A PREFERRED FORMULARY ALTERNATIVE INCLUDING HARVONI OR EPCLUSA WHEN THESE AGENTS ARE CONSIDERED ACCEPTABLE FOR TREATMENT OF THE SPECIFIC GENOTYPE PER AASLD/IDSA GUIDANCE. MUST BE USED CONCURRENTLY WITH RIBAVIRIN UNLESS PATIENT IS TREATMENT NAIVE AND HAS CONTRAINDICATION TO RIBAVIRIN. PATIENT IS NOT CONCURRENTLY TAKING ANY OF THE FOLLOWING (CONTRAINDICATED OR NOT RECOMMENDED BY THE MANUFACTURER): ALFUZOSIN, CARBAMAZEPINE, PHENYTOIN, PHENOBARBITAL, RIFAMPIN, ERGOTAMINE, DIHYDROERGOTAMINE, ERGONOVINE, METHYLERGONOVINE, ETHINYL ESTRADIOL CONTAINING MEDICATIONS (SUCH AS COMBINED ORAL CONTRACEPTIVES, NUVARING, ORTHO EVRA OR XULANE TRANSDERMAL PATCH SYSTEM), LOVASTATIN, SIMVASTATIN, PIMOZIDE, EFAVIRENZ (ATRIPLA, SUSTIVA), REVATIO (SILDENAFIL DOSE OF 20MG AND/OR DOSED THREE TIMES DAILY FOR PAH), TRIAZOLAM, ORAL MIDAZOLAM, LOPINAVIR/RITONAVIR, RILPIVIRINE, SALMETEROL.</p>

OMBITASVIR-PARITAPREVIR-RITONAVIR-DASABUVIR

Products Affected

- Viekira Pak
- Viekira XR

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D. CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT AASLD-IDSA GUIDANCE AND ADDITIONAL CONSIDERATION FOR COVERAGE CONSISTENT WITH FDA LABELING.
Exclusion Criteria	DECOMPENSATED CIRRHOSIS, MODERATE OR SEVERE LIVER IMPAIRMENT (CHILD-PUGH B OR C).
Required Medical Information	HCV RNA LEVEL WITHIN PAST 6 MONTHS.
Age Restrictions	
Prescriber Restrictions	GASTROENTEROLOGIST, INFECTIOUS DISEASE SPECIALIST, PHYSICIAN SPECIALIZING IN THE TREATMENT OF HEPATITIS (HEPATOLOGIST), OR A SPECIALLY TRAINED GROUP SUCH AS ECHO (EXTENSION FOR COMMUNITY HEALTHCARE OUTCOMES) MODEL.
Coverage Duration	CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT AASLD/IDSA GUIDANCE.

PA Criteria	Criteria Details
Other Criteria	<p>CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT AASLD/IDSA GUIDANCE. TRIAL OF A PREFERRED FORMULARY ALTERNATIVE INCLUDING HARVONI OR EPCLUSA WHEN THESE AGENTS ARE CONSIDERED ACCEPTABLE FOR TREATMENT OF THE SPECIFIC GENOTYPE PER AASLD/IDSA GUIDANCE. PATIENT IS NOT CONCURRENTLY TAKING ANY OF THE FOLLOWING: ALFUZOSIN, CARBAMAZEPINE, PHENYTOIN, PHENOBARBITAL, GEMFIBROZIL, RIFAMPIN, ERGOTAMINE, DIHYDROERGOTAMINE, ERGONOVINE, METHYLERGONOVINE, ETHINYL ESTRADIOL CONTAINING MEDICATIONS (SUCH AS COMBINED ORAL CONTRACEPTIVES, NUVARING, ORTHO EVRA OR XULANE TRANSDERMAL PATCH SYSTEM), ST. JOHN'S WORT, LOVASTATIN, SIMVASTATIN, PIMOZIDE, EFAVIRENZ, REVATIO, TRIAZOLAM, ORAL MIDAZOLAM, DARUNAVIR/RITONAVIR, LOPINAVIR/RITONAVIR, RILPIVIRINE, SALMETEROL.</p>

OPIOID DEPENDENCY AGENTS

Products Affected

- buprenorphine HCl sublingual

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	BUPRENORPHINE: 2 DAYS UNLESS PREGNANT (42WKS) OR SERIOUS NALOXONE ALLERGY (24WKS)
Other Criteria	BUPRENORPHINE MONOTHERAPY WILL BE APPROVED FOR THE FOLLOWING PATIENTS: PREGNANCY OR ATTESTATION OF SERIOUS NALOXONE ALLERGY DEFINED AS NALOXONE-INDUCED ANAPHYLAXIS, BRONCHOSPASM, OR ANGIONEUROTIC EDEMA, AND FOR PATIENTS BEING TRANSITIONED DIRECTLY FROM A LONG ACTING OPIOID (I.E. METHADONE, FENTANYL PATCH, OR OTHER ER OPIOIDS). FOR PATIENTS BEING TRANSITIONED DIRECTLY FROM A LONG-ACTING OPIOID APPROVAL FOR BUPRENORPHINE MONOTHERAPY WILL BE LIMITED TO USE DURING INDUCTION THERAPY ONLY (2 DAYS). FOR NEW STARTS: ADDITIONAL CONSIDERATION WILL BE PROVIDED FOR A QUANTITY EXCEPTION FOR INDUCTION DOSING FOR 2 DAYS OF THERAPY.

OSIMERTINIB

Products Affected

- Tagrisso

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	CONCURRENT THERAPY WITH AN EGFR TYROSINE KINASE-INHIBITOR
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	

OXYMETHOLONE

Products Affected

- Anadrol-50

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	CARCINOMA OF THE PROSTATE OR BREAST IN MALE PATIENTS, CARCINOMA OF THE BREAST IN FEMALES WITH HYPERCALCEMIA, WOMEN WHO ARE OR MAY BECOME PREGNANT, NEPHROSIS OR THE NEPHROTIC PHASE OF NEPHRITIS, HYPERSENSITIVITY TO THE DRUG AND SEVERE HEPATIC DYSFUNCTION.
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	

PALBOCICLIB

Products Affected

- Ibrance

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PREVIOUS TRIAL OF OR CONTRAINDICATION TO THE PREFERRED FORMULARY ALTERNATIVE RIBOCICLIB (KISQALI).

PALIVIZUMAB

Products Affected

- Synagis

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	GESTATIONAL AGE
Age Restrictions	LESS THAN 24 MONTHS OF AGE.
Prescriber Restrictions	
Coverage Duration	1 MONTH TO 5 MONTHS. SEE OTHER CRITERIA FOR MORE INFORMATION.
Other Criteria	CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT RECOMMENDATIONS FROM THE AMERICAN ACADEMY OF PEDIATRICS FOR PALIVIZUMAB PROPHYLAXIS FOR RESPIRATORY SYNCYTIAL VIRUS INFECTIONS. INITIAL: APPROVAL WILL BE FOR AT LEAST 1 MONTH AND NO GREATER THAN 5 MONTHS DEPENDENT UPON REMAINING LENGTH OF RESPIRATORY SYNCYTIAL VIRUS (RSV) SEASON. RENEWAL: ADDITIONAL 1 MONTH OF TREATMENT FOR CARDIOPULMONARY BYPASS SURGERY DURING RSV PROPHYLAXIS SEASON.

PANITUMUMAB

Products Affected

- Vectibix

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	

PANOBINOSTAT

Products Affected

- Farydak

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	RENEWAL: PATIENT HAS TOLERATED THE FIRST 8 CYCLES OF THERAPY WITHOUT UNRESOLVED SEVERE OR MEDICALLY SIGNIFICANT TOXICITY.

PARATHYROID HORMONE

Products Affected

- Natpara

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	

PAZOPANIB

Products Affected

- Votrient

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	

PDE5 INHIBITORS FOR PULMONARY ARTERIAL HYPERTENSION

Products Affected

- Adcirca
- sildenafil oral

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	PATIENT CANNOT CONCURRENTLY OR INTERMITTENTLY BE TAKING ORAL ERECTILE DYSFUNCTION AGENTS (E.G. CIALIS, VIAGRA), ANY ORGANIC NITRATES IN ANY FORM, OR GUANYLATE CYCLASE (GC) STIMULATORS (ADEMPAS).
Required Medical Information	DOCUMENTED CONFIRMATORY PULMONARY ARTERIAL HYPERTENSION (PAH) DIAGNOSIS BASED ON RIGHT HEART CATHETERIZATION. PATIENT HAS NYHA-WHO FUNCTIONAL CLASS II-IV SYMPTOMS
Age Restrictions	
Prescriber Restrictions	PRESCRIBED BY OR GIVEN IN CONSULTATION WITH A CARDIOLOGIST OR PULMONOLOGIST
Coverage Duration	INITIAL AND RENEWAL: 12 MONTHS
Other Criteria	INITIAL: MEAN PULMONARY ARTERY PRESSURE (PAP) OF AT LEAST 25 MMHG OR GREATER, PULMONARY CAPILLARY WEDGE PRESSURE (PCWP) OF 15 MMHG OR LESS, PULMONARY VASCULAR RESISTANCE (PVR) GREATER THAN 3 WOOD UNITS. REQUEST FOR ADCIRCA REQUIRE TRIAL OR CONTRAINDICATION TO REVATIO. RENEWAL: PATIENT SHOWS IMPROVEMENT FROM BASELINE IN THE 6-MINUTE WALK DISTANCE OR PATIENT HAS A STABLE 6-MINUTE WALK DISTANCE WITH A STABLE/ IMPROVED WHO FUNCTIONAL CLASS. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION.

PDE5 INHIBITORS FOR PULMONARY ARTERIAL HYPERTENSION - IV

Products Affected

- sildenafil intravenous

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	PATIENT CANNOT CONCURRENTLY OR INTERMITTENTLY BE TAKING ORAL ERECTILE DYSFUNCTION AGENTS (E.G. CIALIS, VIAGRA), ANY ORGANIC NITRATES IN ANY FORM, OR GUANYLATE CYCLASE (GC) STIMULATORS (ADEMPAS).
Required Medical Information	DOCUMENTED CONFIRMATORY PULMONARY ARTERIAL HYPERTENSION (PAH) DIAGNOSIS BASED ON RIGHT HEART CATHETERIZATION. PATIENT HAS NYHA-WHO FUNCTIONAL CLASS II-IV SYMPTOMS
Age Restrictions	
Prescriber Restrictions	PRESCRIBED BY OR GIVEN IN CONSULTATION WITH A CARDIOLOGIST OR PULMONOLOGIST
Coverage Duration	INITIAL AND RENEWAL: 12 MONTHS
Other Criteria	INITIAL: MEAN PULMONARY ARTERY PRESSURE (PAP) OF AT LEAST 25 MMHG OR GREATER, PULMONARY CAPILLARY WEDGE PRESSURE (PCWP) OF 15 MMHG OR LESS, PULMONARY VASCULAR RESISTANCE (PVR) GREATER THAN 3 WOOD UNITS. RENEWAL: PATIENT SHOWS IMPROVEMENT FROM BASELINE IN THE 6-MINUTE WALK DISTANCE OR PATIENT HAS A STABLE 6-MINUTE WALK DISTANCE WITH A STABLE/ IMPROVED WHO FUNCTIONAL CLASS.

PEG-INTERFERON ALFA-2B-SYLATRON

Products Affected

- Sylatron

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	OVERALL DURATION OF THERAPY LIMITED TO 5 YEARS.

PEMBROLIZUMAB

Products Affected

- Keytruda intravenous recon soln
- Keytruda intravenous solution

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	

PENICILLAMINE

Products Affected

- Cuprimine

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	RHEUMATOID ARTHRITIS: HISTORY OR OTHER EVIDENCE OF RENAL INSUFFICIENCY
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	WILSON'S DISEASE: PRESCRIBED BY OR GIVEN IN CONSULTATION WITH A HEPATOLOGIST. CYSTINURIA: PRESCRIBED BY OR GIVEN IN CONSULTATION WITH A NEPHROLOGIST. RHEUMATOID ARTHRITIS: PRESCRIBED BY OR GIVEN IN CONSULTATION WITH A RHEUMATOLOGIST.
Coverage Duration	12 MONTHS
Other Criteria	WILSON'S DISEASE: GENETIC TESTING FOR ATP7B MUTATIONS. CYSTINURIA: DIAGNOSIS REQUIRES THE PRESENCE OF NEPHROLITHIASIS AND 1 OR MORE OF THE FOLLOWING: STONE ANALYSIS SHOWING PRESENCE OF CYSTEINE, IDENTIFICATION OF PATHOGNOMONIC HEXAGONAL CYSTINE CRYSTALS ON URINALYSIS, POSITIVE FAMILY HISTORY OF CYSTINURIA WITH POSITIVE CYANIDE-NITROPRUSSIDE SCREEN. REQUESTS FOR CUPRIMINE FOR THE TREATMENT OF WILSONS DISEASE, CYSTINURIA, AND RHEUMATOID ARTHRITIS REQUIRE A PREVIOUS TRIAL OF OR CONTRAINDICATION TO DEPEN.

PENICILLAMINE-DEPEN

Products Affected

- Depen Titratabs

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	RHEUMATOID ARTHRITIS; HISTORY OR OTHER EVIDENCE OF RENAL INSUFFICIENCY
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	WILSON'S DISEASE: PRESCRIBED BY OR GIVEN IN CONSULTATION WITH A HEPATOLOGIST. CYSTINURIA: PRESCRIBED BY OR GIVEN IN CONSULTATION WITH A NEPHROLOGIST. RHEUMATOID ARTHRITIS: PRESCRIBED BY OR GIVEN IN CONSULTATION WITH A RHEUMATOLOGIST.
Coverage Duration	12 MONTHS
Other Criteria	WILSON'S DISEASE: GENETIC TESTING FOR ATP7B MUTATIONS. CYSTINURIA: DIAGNOSIS REQUIRES THE PRESENCE OF NEPHROLITHIASIS AND 1 OR MORE OF THE FOLLOWING: STONE ANALYSIS SHOWING PRESENCE OF CYSTEINE, IDENTIFICATION OF PATHOGNOMONIC HEXAGONAL CYSTINE CRYSTALS ON URINALYSIS, POSITIVE FAMILY HISTORY OF CYSTINURIA WITH POSITIVE CYANIDE-NITROPRUSSIDE SCREEN.

PERTUZUMAB

Products Affected

- Perjeta

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	INITIAL: 4 MONTHS. RENEWAL: 12 MONTHS
Other Criteria	

PIMAVANSERIN

Products Affected

- Nuplazid

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	18 YEARS OR OLDER
Prescriber Restrictions	PRESCRIBED BY OR GIVEN IN CONSULTATION WITH A NEUROLOGIST, GERIATRICIAN, OR A BEHAVIORAL HEALTH SPECIALIST (SUCH AS A PSYCHIATRIST).
Coverage Duration	INITIAL 12 MONTHS. RENEWAL 12 MONTHS.
Other Criteria	RENEWAL REQUIRES THAT THE PATIENT HAS EXPERIENCED AN IMPROVEMENT IN PSYCHOSIS SYMPTOMS FROM BASELINE AND DEMONSTRATES A CONTINUED NEED FOR TREATMENT.

PIRFENIDONE

Products Affected

- Esbriet oral capsule
- Esbriet oral tablet 267 mg, 801 mg

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	PATIENTS WITH KNOWN CAUSES OF INTERSTITIAL LUNG DISEASE (E.G., CONNECTIVE TISSUE DISEASE, DRUG TOXICITY, ASBESTOS OR BERYLLIUM EXPOSURE, HYPERSENSITIVITY PNEUMONITIS, SYSTEMIC SCLEROSIS, RHEUMATOID ARTHRITIS, RADIATION, SARCOIDOSIS, BRONCHIOLITIS OBLITERANS ORGANIZING PNEUMONIA, HUMAN IMMUNODEFICIENCY VIRUS (HIV) INFECTION, VIRAL HEPATITIS, AND CANCER). NOT APPROVED IF THE PATIENT HAS NOT OBTAINED LIVER FUNCTION TESTS.
Required Medical Information	PATIENT WITH USUAL INTERSTITIAL PNEUMONIA (UIP) PATTERN AS EVIDENCED BY HIGH-RESOLUTION COMPUTED TOMOGRAPHY (HRCT) ALONE OR VIA A COMBINATION OF SURGICAL LUNG BIOPSY AND HRCT
Age Restrictions	
Prescriber Restrictions	PRESCRIBED BY OR GIVEN IN CONSULTATION WITH A PULMONOLOGIST
Coverage Duration	12 MONTHS
Other Criteria	PATIENT HAS A PREDICTED FORCED VITAL CAPACITY (FVC) OF AT LEAST 50%.

POMALIDOMIDE

Products Affected

- Pomalyst

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	

PONATINIB

Products Affected

- Iclusig oral tablet 15 mg, 45 mg

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	

PRAMLINTIDE

Products Affected

- SymlinPen 120
- SymlinPen 60

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	TYPE I OR TYPE II DIABETES: REQUIRING INSULIN OR CONTINUOUS INSULIN INFUSION (INSULIN PUMP) FOR GLYCEMIC CONTROL
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	

PYRIMETHAMINE

Products Affected

- Daraprim

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D, ADDITIONAL CONSIDERATION FOR CHRONIC MAINTENANCE THERAPY FOR TOXOPLASMOSIS AND TOXOPLASMOSIS PROPHYLAXIS.
Exclusion Criteria	
Required Medical Information	MALARIA: PLASMODIA SUSCEPTIBLE TESTING. TOXOPLASMOSIS:CD4 LEVEL
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	ACUTE MALARIA AND CHEMOPROPHYLAXIS: INITIAL: 3 MONTHS. RENEWAL: 12 MONTHS. SEE OTHER CRITERIA FIELD

PA Criteria	Criteria Details
Other Criteria	<p>INITIAL: ACUTE MALARIA TREATMENT AND MALARIA CHEMOPROPHYLAXIS REQUIRES THAT THE PATIENT HAS MALARIA SUSCEPTIBLE TO PYRIMETHAMINE AND A PREVIOUS TRIAL OF PLAQUENIL (HYDROXYCHLOROQUINE SULFATE) AND MALARONE (ATOVAQUONE/PROGUANIL) (UNLESS THESE REGIMENS ARE RESISTANT IN THE SPECIFIC REGION AS INDICATED BY REGIONAL PLASMODIA SUSCEPTIBILITY). PRIMARY PROPHYLAXIS OF TOXOPLASMOSIS IN PATIENTS WITH HIV REQUIRES PREVIOUS TRIAL OF OR CONTRAINDICATION TO BACTRIM (SMX/TMP). RENEWAL: CONTINUATION OF TREATMENT FOLLOWING ACUTE MALARIA REQUIRES PREVIOUS INFECTION WITH MALARIA SUSCEPTIBLE TO PYRIMETHAMINE WITH SUBSEQUENT CLINICAL CURE (ELIMINATION OF MALARIA SYMPTOMS DEFINED AS CHILLS, FEVER, SWEATS, GENERAL MALAISE) FOLLOWED BY SYMPTOMS OF RELAPSE. CONTINUATION OF MALARIA CHEMOPROPHYLAXIS REQUIRES THE PATIENT WILL BE TRAVELING TO OR RESIDING IN AN AREA WHERE PLASMODIA SUSCEPTIBLE TO PYRIMETHAMINE EXISTS (MALARIA MUST BE SENSITIVE TO PYRIMETHAMINE).CONTINUED TREATMENT OF TOXOPLASMOSIS REQUIRES ONE OF THE FOLLOWING: 1) PERSISTENT CLINICAL DISEASE (HEADACHE, NEUROLOGICAL SYMPTOMS, OR FEVER) AND PERSISTENT RADIOGRAPHIC DISEASE (ONE OR MORE MASS LESIONS ON BRAIN IMAGING) OR 2) CD4 COUNT LESS THAN 200 CELLS/MM3 AND CURRENT ANTI-RETROVIRAL THERAPY IF HIV POSITIVE. CONTINUATION OF PRIMARY PROPHYLAXIS FOR TOXOPLASMOSIS WITH HIV REQUIRES CD4 COUNT LESS THAN 200 CELLS/MM3 AND CURRENT ANTI RETROVIRAL THERAPY. TOXOPLASMOSIS: INITIAL: 8 WEEKS. RENEWAL: 6 MONTHS. PRIMARY PROPHYLAXIS OF TOXOPLASMOSIS: INITIAL AND RENEWAL IS 12 MONTHS.</p>

QUININE SULFATE

Products Affected

- quinine sulfate

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	

RAMUCIRUMAB

Products Affected

- Cyramza

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	

REGORAFENIB

Products Affected

- Stivarga

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	FOR COLORECTAL CANCER: TRIAL OF OR CONTRAINDICATION TO AN ANTI-VEGF THERAPY SUCH AS AVASTIN OR ZALTRAP AND A FLUOROPYRIMIDINE-, OXALIPLATIN- AND IRINOTECAN-BASED CHEMOTHERAPY SUCH AS FOLFOX, FOLFOXIRI, FOLFIRI, CAPEOX, INFUSIONAL 5-FU/LV OR CAPECITABINE. IF APPLICABLE, A TRIAL OF OR CONTRAINDICATION TO AN ANTI-EGFR THERAPY SUCH AS ERBITUX OR VECTIBIX IS ALSO REQUIRED FOR KRAS WILD TYPE COLORECTAL CANCER. FOR GIST, A TRIAL OF OR CONTRAINDICATION TO GLEEVEC AND SUTENT IS REQUIRED.

RESLIZUMAB

Products Affected

- Cinqair

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	CONCURRENT USE OF XOLAIR
Required Medical Information	BLOOD EOSINOPHIL LEVEL GREATER THAN OR EQUAL TO 400 CELLS/MCL WITHIN THE LAST 6 MONTHS
Age Restrictions	
Prescriber Restrictions	PRESCRIBED BY OR GIVEN IN CONSULTATION WITH A PHYSICIAN SPECIALIZING IN PULMONARY MEDICINE
Coverage Duration	INITIAL 24 WEEKS. RENEWAL 12 MONTHS
Other Criteria	INITIAL THERAPY: PATIENT CURRENTLY TREATED WITH A MAXIMALLY TOLERATED DOSE OF INHALED CORTICOSTEROIDS. RENEWAL REQUIRES DOCUMENTATION THAT THE PATIENT HAS EXPERIENCED AT LEAST A 25 PERCENT REDUCTION IN ASTHMA EXACERBATIONS (FOR EXAMPLE: HOSPITALIZATIONS, URGENT OR EMERGENT CARE VISITS, USE OF RESCUE MEDICATIONS, ETC.) FROM BASELINE.

RIBOCICLIB

Products Affected

- Kisqali
- Kisqali Femara Co-Pack oral tablet 200 mg/day(200 mg x 1)-2.5 mg, 400 mg/day(200 mg x 2)-2.5 mg, 600 mg/day(200 mg x 3)-2.5 mg

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	

RIFAXIMIN

Products Affected

- Xifaxan oral tablet 200 mg, 550 mg

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	TRAVELERS' DIARRHEA:1 FILL IN 1MONTH.HEPATIC ENCEPHALOPATHY:12 MO.IBS-D:INITIAL:12 WKS.RENEWAL:12 MO
Other Criteria	FOR RIFAXIMIN 550 MG TABLETS ONLY: INITIAL: HEPATIC ENCEPHALOPATHY (HE): PREVIOUS TRIAL OF OR CONTRAINDICATION TO LACTULOSE OR CONCURRENT LACTULOSE THERAPY. IRRITABLE BOWEL SYNDROME WITH DIARRHEA (IBS-D): PREVIOUS TRIAL OF OR CONTRAINDICATION TO DICYCLOMINE. RENEWAL FOR IBS-D REQUIRES THAT AT LEAST 10 WEEKS HAVE PASSED SINCE THE LAST TREATMENT COURSE OF RIFAXIMIN, THE PATIENT HAS EXPERIENCED AT LEAST A 30% DECREASE IN ABDOMINAL PAIN (ON A 0-10 POINT PAIN SCALE) AND A 50% REDUCTION IN THE NUMBER OF DAYS PER WEEK WITH A STOOL CONSISTENCY OF MUSHY STOOL (BRISTOL STOOL SCALE TYPE 6) OR ENTIRELY LIQUID STOOL (BRISTOL STOOL SCALE TYPE 7).

RIOCIGUAT

Products Affected

- Adempas

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	INITIAL FOR PAH: PATIENT IS NOT CONCURRENTLY TAKING NITRATES OR NITRIC OXIDE DONORS (E.G. AMYL NITRATE), PHOSPHODIESTERASE INHIBITORS (E.G. SILDENAFIL, TADALAFIL, OR VARDENAFIL), OR NON-SPECIFIC PDE INHIBITORS (E.G. DIPYRIDAMOLE, THEOPHYLLINE). INITIAL FOR CTEPH: PATIENT IS NOT A CANDIDATE FOR SURGERY OR HAS INOPERABLE CTEPH. PERSISTENT OR RECURRENT DISEASE AFTER SURGICAL TREATMENT. PATIENT IS NOT CONCURRENTLY OR INTERMITTENTLY TAKING NITRATES, NITRIC OXIDE DONORS OR ANY PDE INHIBITORS (E.G.VIAGRA, CIALIS, DIPYRIDAMOLE).
Required Medical Information	DOCUMENTED CONFIRMATORY PULMONARY ARTERIAL HYPERTENSION (PAH) DIAGNOSIS BASED ON RIGHT HEART CATHETERIZATION. PATIENT HAS NYHA-WHO FUNCTIONAL CLASS II-IV SYMPTOMS. DIAGNOSIS OF PERSISTENT/RECURRENT CHRONIC THROMBOEMBOLIC PULMONARY HYPERTENSION (CTEPH) WHO GROUP 4. PATIENT HAS NYHA-WHO FUNCTIONAL CLASS II-IV SYMPTOMS.
Age Restrictions	
Prescriber Restrictions	PRESCRIBED BY OR IN CONSULTATION WITH A CARDIOLOGIST OR PULMONOLOGIST.
Coverage Duration	INITIAL AND RENEWAL: 12 MONTHS

PA Criteria	Criteria Details
Other Criteria	<p>INITIAL FOR PAH: MEAN PULMONARY ARTERY PRESSURE (PAP) OF AT LEAST 25 MMHG OR GREATER, PULMONARY CAPILLARY WEDGE PRESSURE (PCWP) OF 15 MMHG OR LESS, PULMONARY VASCULAR RESISTANCE (PVR) GREATER THAN 3 WOOD UNITS. PREVIOUS TRIAL OF OR CONTRAINDICATION TO A PHOSPHODIESTERASE-5 (PDE-5) INHIBITOR, SUCH AS REVATIO OR ADCIRCA.</p> <p>RENEWAL FOR PAH AND CTEPH: PATIENT SHOW IMPROVEMENT FROM BASELINE IN THE 6-MINUTE WALK DISTANCE OR PATIENT HAS A STABLE 6-MINUTE WALK DISTANCE WITH A STABLE/ IMPROVED WHO FUNCTIONAL CLASS.</p>

RITUXIMAB

Products Affected

- Rituxan

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	RENEWAL: PHYSICIAN ATTESTATION OF IMPROVEMENT.
Age Restrictions	
Prescriber Restrictions	RHEUMATOID ARTHRITIS: PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. NHL, CLL: ONCOLOGIST.
Coverage Duration	RA: INITIAL: 6 MO. RENEWAL: 12 MONTHS. NHL: 1 YEAR. CLL: 6 MO. WG, MPA: 3 MONTH.
Other Criteria	INITIAL: RHEUMATOID ARTHRITIS: PREVIOUS TRIAL OF HUMIRA FOLLOWED BY ONE OF THE FOLLOWING PREFERRED AGENTS: ORENCIA, XELJANZ, CIMZIA OR ACTEMRA.

RITUXIMAB SQ

Products Affected

- Rituxan Hycela

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	THE PATIENT HAS RECEIVED OR WILL RECEIVE AT LEAST ONE FULL DOSE OF A RITUXIMAB PRODUCT BY INTRAVENOUS INFUSION PRIOR TO INITIATION OF RITUXIMAB AND HYALURONIDASE.

ROMIPLOSTIM

Products Affected

- Nplate

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	INITIAL: ADEQUATE RESPONSE TO CORTICOSTEROIDS, IMMUNOGLOBULINS, OR SPLENECTOMY. RENEWAL: NO CLINICAL RESPONSE AS DEFINED BY AN INCREASE IN PLATELET COUNT OF GREATER THAN OR EQUAL TO $50 \times 10^9/L$ AT THE MAX DOSE OF 10 MCG/KG PER DAY FOR 4 WEEKS.
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	INITL: 2 MO., RENEW: IF NO RESPONSE AFTER INITIAL APPROVAL: 1 MO. AT MAX DOSE. IF RESPONSE: 12 MO.
Other Criteria	

RUCAPARIB

Products Affected

- Rubraca

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	

RUXOLITINIB

Products Affected

- Jakafi

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	RENEWAL: IMPROVEMENT OR MAINTENANCE OF SYMPTOM IMPROVEMENT SUCH AS A 50% OR GREATER REDUCTION IN TOTAL SYMPTOM SCORE ON THE MODIFIED MYELOFIBROSIS SYMPTOM ASSESSMENT FORM (MFSAF) V2.0 OR 50% OR GREATER REDUCTION IN PALPABLE SPLEEN LENGTH, OR REDUCTION OF 35% OR GREATER FROM BASELINE SPLEEN VOLUME AFTER 6 MONTHS OF THERAPY.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS
Other Criteria	

SAFINAMIDE MESYLATE

Products Affected

- Xadago

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	

SARILUMAB

Products Affected

- Kevzara

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	RENEWAL: PHYSICIAN ATTESTATION OF IMPROVEMENT.
Age Restrictions	
Prescriber Restrictions	PRESCRIBED BY OR GIVEN IN CONSULTATION WITH A RHEUMATOLOGIST.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS
Other Criteria	INITIAL: PREVIOUS TRIAL OF HUMIRA AND ONE OF THE FOLLOWING PREFERRED AGENTS: ACTEMRA, CIMZIA, ORENCIA, OR XELJANZ.

SEBELIPASE ALFA

Products Affected

- Kanuma

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	BLOOD TEST OR DRIED BLOOD SPOT TEST INDICATING LOW OR ABSENT LYSOSOMAL ACID LIPASE DEFICIENCY (LAL) ENZYME ACTIVITY, OR A GENETIC TEST INDICATING THE PRESENCE OF ALTERED LIPA GENE(S).
Age Restrictions	
Prescriber Restrictions	PRESCRIBED BY OR GIVEN IN CONSULTATION WITH AN ENDOCRINOLOGIST, HEPATOLOGIST, GASTROENTEROLOGIST, MEDICAL GENETICIST, LIPIDOLOGIST, OR A METABOLIC SPECIALIST.
Coverage Duration	LAL INITIAL 6 OR 12 MONTHS, SEE OTHER CRITERIA. RENEWAL: 12 MONTHS

PA Criteria	Criteria Details
Other Criteria	<p>INITIAL: DIAGNOSIS OF LYSOSOMAL ACID LIPASE (LAL) DEFICIENCY, AS CONFIRMED BY THE PRESENCE OF CLINICAL FEATURES (E.G., HEPATOMEGALY, ELEVATED SERUM TRANSAMINASES, DYSLIPIDEMIA, SPLENOMEGALY) PLUS ANY OF THE FOLLOWING: A BLOOD TEST INDICATING LOW OR ABSENT LEVELS OF LAL ENZYME ACTIVITY, A DRIED BLOOD SPOT TEST INDICATING LOW OR ABSENT LAL ENZYME ACTIVITY, OR A GENETIC TEST INDICATING THE BI-ALLELIC PRESENCE OF ALTERED LIPA GENE(S).</p> <p>RENEWAL:DIAGNOSIS OF LYSOSOMAL ACID LIPASE (LAL) DEFICIENCY PRESENTING AFTER THE FIRST 6 MONTHS OF LIFE AND NOT CONSIDERED RAPIDLY PROGRESSIVE REQUIRES DOCUMENTED IMPROVEMENT IN ANY ONE OF THE FOLLOWING CLINICAL PARAMETERS ASSOCIATED WITH LYSOSOMAL ACID LIPASE (LAL) DEFICIENCY DURING THE PAST 6 MONTHS: A RELATIVE REDUCTION FROM BASELINE IN ANY ONE OF THE FOLLOWING LIPID LEVELS (LDL-C, NON-HDL-C, OR TRIGLYCERIDES), NORMALIZATION OF ASPARTATE AMINOTRANSFERASE (AST) BASED ON AGE- AND GENDER-SPECIFIC NORMAL RANGES, A DECREASE IN LIVER FAT CONTENT COMPARED TO BASELINE ASSESSED BY ABDOMINAL IMAGING (E.G., MULTI-ECHO GRADIENT ECHO [MEGE] MRI). DIAGNOSIS OF RAPIDLY PROGRESSIVE LYSOSOMAL ACID LIPASE (LAL) DEFICIENCY PRESENTING WITHIN THE FIRST 6 MONTHS OF LIFE: 12 MONTHS. A DIAGNOSIS OF LYSOSOMAL ACID LIPASE (LAL) DEFICIENCY PRESENTING AFTER THE FIRST 6 MONTHS OF LIFE AND NOT CONSIDERED RAPIDLY PROGRESSIVE: INITIAL: 6 MONTHS</p>

SECUKINUMAB

Products Affected

- Cosentyx (2 Syringes)
- Cosentyx Pen (2 Pens)

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	PLAQUE PSORIASIS (PSO): MODERATE TO SEVERE PLAQUE PSORIASIS INVOLVING GREATER THAN OR EQUAL TO 5 PERCENT BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, OR GENITAL AREA. RENEWAL: PHYSICIAN ATTESTATION OF IMPROVEMENT.
Age Restrictions	
Prescriber Restrictions	PLAQUE PSORIASIS (PSO): PRESCRIBED BY OR GIVEN IN CONSULTATION WITH A DERMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR GIVEN IN CONSULTATION WITH A RHEUMATOLOGIST OR A DERMATOLOGIST. ANKYLOSING SPONDYLITIS: PRESCRIBED BY OR GIVEN IN CONSULTATION WITH A RHEUMATOLOGIST.
Coverage Duration	INITIAL: 4 MONTHS. RENEWAL: 12 MONTHS FOR ALL DIAGNOSES
Other Criteria	INITIAL: PLAQUE PSORIASIS (PSO): PREVIOUS TRIAL OF HUMIRA AND ONE CONVENTIONAL THERAPY SUCH AS PUVA (PHOTOTHERAPY ULTRAVIOLET LIGHT A), UVB (ULTRAVIOLET LIGHT B), TOPICAL CORTICOSTEROIDS, CALCIPOTRIENE, ACITRETIN, METHOTREXATE, OR CYCLOSPORINE. PSORIATIC ARTHRITIS (PSA): PREVIOUS TRIAL OF HUMIRA AND ONE DMARD (DISEASE-MODIFYING ANTIRHEUMATIC DRUG) SUCH AS METHOTREXATE, LEFLUNOMIDE, HYDROXYCHLOROQUINE, OR SULFASALAZINE. ANKYLOSING SPONDYLITIS (AS): PREVIOUS TRIAL OF HUMIRA.

SELEXIPAG

Products Affected

- Upravi oral tablet 1,000 mcg, 1,200 mcg, 1,400 mcg, 1,600 mcg, 200 mcg, 400 mcg, 600 mcg, 800 mcg
- Upravi oral tablets, dose pack

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	DOCUMENTED CONFIRMATORY PULMONARY ARTERIAL HYPERTENSION (PAH) DIAGNOSIS BASED ON RIGHT HEART CATHETERIZATION. PATIENT HAS NYHA-WHO FUNCTIONAL CLASS II-IV SYMPTOMS.
Age Restrictions	
Prescriber Restrictions	PRESCRIBED BY OR GIVEN IN CONSULTATION WITH A CARDIOLOGIST OR PULMONOLOGIST
Coverage Duration	INITIAL AND RENEWAL: 12 MONTHS
Other Criteria	INITIAL: MEAN PULMONARY ARTERY PRESSURE (PAP) OF AT LEAST 25 MMHG OR GREATER, PULMONARY CAPILLARY WEDGE PRESSURE (PCWP) OF 15 MMHG OR LESS, PULMONARY VASCULAR RESISTANCE (PVR) GREATER THAN 3 WOOD UNITS. RENEWAL: PATIENT SHOWS IMPROVEMENT FROM BASELINE IN THE 6-MINUTE WALK DISTANCE OR PATIENT HAS A STABLE 6-MINUTE WALK DISTANCE WITH A STABLE/ IMPROVED WHO FUNCTIONAL CLASS.

SILTUXIMAB

Products Affected

- Sylvant

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	

SIMEPREVIR

Products Affected

- Olysio

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D. CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT AASLD-IDSA GUIDANCE AND ADDITIONAL CONSIDERATION FOR COVERAGE CONSISTENT WITH FDA LABELING.
Exclusion Criteria	
Required Medical Information	HCV RNA LEVEL WITHIN PAST 6 MONTHS. FOR ALL GENOTYPE 1A: NS3 80K POLYMORPHISM LAB TEST AT BASELINE.
Age Restrictions	
Prescriber Restrictions	GASTROENTEROLOGIST, INFECTIOUS DISEASE SPECIALIST, PHYSICIAN SPECIALIZING IN THE TREATMENT OF HEPATITIS (FOR EXAMPLE HEPATOLOGIST), OR A SPECIALLY TRAINED GROUP SUCH AS ECHO (EXTENSION FOR COMMUNITY HEALTHCARE OUTCOMES) MODEL.
Coverage Duration	CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT AASLD-IDSA GUIDANCE.

PA Criteria	Criteria Details
Other Criteria	<p>CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT AASLD-IDSa GUIDANCE. TRIAL OF A PREFERRED FORMULARY ALTERNATIVE INCLUDING HARVONI OR EPCLUSA WHEN THESE AGENTS ARE CONSIDERED ACCEPTABLE FOR TREATMENT OF THE SPECIFIC GENOTYPE PER AASLD/IDSa GUIDANCE. PATIENT MUST NOT BE TAKING ANY OF THE FOLLOWING INTERACTING MEDICATIONS: CARBAMAZEPINE, PHENYTOIN, PHENOBARBITAL, OXCARBAZEPINE, RIFAMPIN, RIFABUTIN, RIFAPENTINE, ERYTHROMYCIN (DOES NOT INCLUDE TOPICAL FORMULATIONS), CLARITHROMYCIN, TELITHROMYCIN, ITRACONAZOLE, KETOCONAZOLE, POSACONAZOLE, FLUCONAZOLE (DOES NOT INCLUDE TOPICAL FORMULATIONS), VORICONAZOLE, DEXAMETHASONE, CISAPRIDE, CYCLOSPORINE, ROSUVASTATIN DOSE ABOVE 10MG, ATORVASTATIN DOSE ABOVE 40MG, OR ANY OF THE FOLLOWING HIV MEDICATIONS: COBICISTAT-CONTAINING MEDS (E.G., STRIBILD), ANY HIV PROTEASE INHIBITOR (ATAZANAVIR, FOSAMPRENAVIR, LOPINAVIR, INDINAVIR, NELFINAVIR, SAQUINAVIR, OR TIPRANAVIR) RITONAVIR, DARUNAVIR/RITONAVIR, DELAVIRDINE, ETRAVIRINE, NEVIRAPINE, EFAVIRENZ). PATIENT MUST ALSO NOT BE TAKING AMIODARONE IF ON COMBINATION REGIMEN OF SOVALDI AND OLYSIO.</p>

SOFOSBUVIR

Products Affected

- Sovaldi

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D. CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT AASLD-IDSA GUIDANCE AND ADDITIONAL CONSIDERATION FOR COVERAGE CONSISTENT WITH FDA LABELING.
Exclusion Criteria	PATIENT WITH END STAGE RENAL DISEASE OR REQUIRES DIALYSIS.
Required Medical Information	FOR ALL GENOTYPE 1 PATIENTS USING OLYSIO AND SOVALDI AND HAVE GENOTYPE 1A: NS3 80K POLYMORPHISM LAB TEST AT BASELINE.
Age Restrictions	
Prescriber Restrictions	PRESCRIBED BY OR GIVEN IN CONSULTATION WITH A GASTROENTEROLOGIST, INFECTIOUS DISEASE SPECIALIST, PHYSICIAN SPECIALIZING IN THE TREATMENT OF HEPATITIS (HEPATOLOGIST), OR A SPECIALLY TRAINED GROUP SUCH AS ECHO (EXTENSION FOR COMMUNITY HEALTHCARE OUTCOMES) MODEL
Coverage Duration	CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT AASLD/IDSA GUIDANCE

PA Criteria	Criteria Details
Other Criteria	<p>CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT AASLD/IDSA GUIDANCE. TRIAL OF A PREFERRED FORMULARY ALTERNATIVE INCLUDING HARVONI OR EPCLUSA WHEN THESE AGENTS ARE CONSIDERED ACCEPTABLE FOR TREATMENT OF THE SPECIFIC GENOTYPE PER AASLD/IDSA GUIDANCE. FOR PATIENTS ON SOVALDI PLUS DAKLINZA REGIMENS THERE WILL BE NO APPROVALS FOR CONCURRENT USE OF ANY OF THESE (CONTRAINDICATED OR NOT RECOMMENDED BY THE MANUFACTURER) MEDICATIONS: AMIODARONE, CARBAMAZEPINE, PHENYTOIN, OR RIFAMPIN. REQUESTS FOR SOVALDI IN COMBINATION WITH DAKLINZA OR OLYSIO WILL REQUIRE THAT THE PATIENT ALSO MEETS ALL CRITERIA FOR THE RESPECTIVE AGENT USED (DAKLINZA OR OLYSIO).</p>

SOFOSBUVIR/VELPATASVIR

Products Affected

- Epclusa

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D. CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT AASLD-IDSA GUIDANCE.
Exclusion Criteria	
Required Medical Information	HCV RNA LEVEL.
Age Restrictions	18 YEARS OF AGE AND OLDER.
Prescriber Restrictions	PRESCRIBED BY OR GIVEN IN CONSULTATION WITH: GASTROENTEROLOGIST, INFECTIOUS DISEASE SPECIALIST, PHYSICIAN SPECIALIZING IN THE TREATMENT OF HEPATITIS (HEPATOLOGIST), OR A SPECIALLY TRAINED GROUP SUCH AS ECHO (EXTENSION FOR COMMUNITY HEALTHCARE OUTCOMES) MODEL.
Coverage Duration	CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT AASLD/IDSA GUIDANCE.
Other Criteria	CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT AASLD/IDSA GUIDANCE. HCV RNA LEVEL WITHIN PAST 6 MONTHS. PATIENT IS NOT CONCURRENTLY TAKING ANY OF THE FOLLOWING MEDICATIONS NOT RECOMMENDED BY THE MANUFACTURER: AMIODARONE, CARBAMAZEPINE, PHENYTOIN, PHENOBARBITAL, OXCARBAZEPINE, RIFAMPIN, RIFABUTIN, RIFAPENTINE, HIV REGIMEN THAT CONTAINS EFAVIRENZ, ROSUVASTATIN AT DOSES ABOVE 10MG, TIPRANA VIR/RITONAVIR OR TOPOTECAN. PATIENT MUST NOT HAVE SEVERE RENAL IMPAIRMENT, ESRD OR ON HEMODIALYSIS. RIBAVIRIN USE REQUIRED FOR PATIENTS WITH DECOMPENSATED CIRRHOSIS.

SOMATROPIN - GROWTH HORMONE

Products Affected

- Humatrope
- Omnitrope
- Saizen
- Saizen click.easy
- Zomacton

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	ATHLETIC ENHANCEMENT, ANTI-AGING PURPOSES, GROWTH FAILURE WITH CLOSED EPIPHYSES.
Required Medical Information	INDUCTION - PATIENT'S HEIGHT AT LEAST 2 STANDARD DEVIATIONS (SD) BELOW THE MEAN HEIGHT FOR NORMAL CHILDREN OF THE SAME AGE AND GENDER. RENEWAL: GROWTH VELOCITY AND/OR TARGET HEIGHT.
Age Restrictions	
Prescriber Restrictions	PRESCRIBED BY OR GIVEN IN CONSULTATION WITH: ENDOCRINOLOGIST.
Coverage Duration	12 MONTHS
Other Criteria	INITIAL: PREVIOUS TRIAL OF PREFERRED FORMULARY ALTERNATIVES NORDITROPIN AND GENOTROPIN PER FDA INDICATION. RENEWAL: PHYSICIAN ATTESTATION OF IMPROVEMENT. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION.

SOMATROPIN - SEROSTIM

Products Affected

- Serostim subcutaneous recon soln 4 mg, 5 mg, 6 mg

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	ATHLETIC ENHANCEMENT, ANTI-AGING PURPOSES
Required Medical Information	HIV/WASTING: MEETS CRITERIA OF WEIGHT LOSS: 10% UNINTENTIONAL WEIGHT LOSS OVER 12 MONTHS, OR 7.5% OVER 6 MONTHS, OR 5% BODY CELL MASS (BCM) LOSS WITHIN 6 MONTHS, OR A BCM LESS THAN 35% (MEN) OF TOTAL BODY WEIGHT AND A BODY MASS INDEX (BMI) LESS THAN 27 KG PER METER SQUARED, OR BCM LESS THAN 23% (WOMEN) OF TOTAL BODY WEIGHT AND A BODY MASS INDEX (BMI) LESS THAN 27 KG PER METER SQUARED, OR BMI LESS THAN 20 KG PER METER SQUARED.
Age Restrictions	
Prescriber Restrictions	PRESCRIBED BY OR GIVEN IN CONSULTATION WITH A GASTROENTEROLOGIST, NUTRITIONAL SUPPORT SPECIALIST (SBS), OR INFECTIOUS DISEASE SPECIALIST
Coverage Duration	3 MONTHS
Other Criteria	HIV/WASTING: CURRENTLY ON ANTIRETROVIRAL THERAPY. IF CURRENTLY ON GROWTH HORMONE, PATIENT HAS SHOWN CLINICAL BENEFIT IN MUSCLE MASS AND WEIGHT OR IF NOT ON GROWTH HORMONE, PATIENT HAS HAD INADEQUATE RESPONSE TO PREVIOUS THERAPY. (I.E. EXERCISE TRAINING, NUTRITIONAL SUPPLEMENTS, APPETITE STIMULANTS OR ANABOLIC STEROIDS). THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION.

SOMATROPIN - ZORBTIVE

Products Affected

- Zorbtive

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	ATHLETIC ENHANCEMENT, ANTI-AGING PURPOSES
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	PRESCRIBED BY OR GIVEN IN CONSULTATION WITH A GASTROENTEROLOGIST
Coverage Duration	SHORT BOWEL: 4 WEEKS ONCE
Other Criteria	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION.

SOMATROPIN-NORDITROPIN AND GENOTROPIN

Products Affected

- Genotropin
- Genotropin MiniQuick
- Norditropin FlexPro

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	ATHLETIC ENHANCEMENT, ANTI-AGING PURPOSES, GROWTH FAILURE WITH CLOSED EPIPHYSES.
Required Medical Information	INDUCTION - PATIENT'S HEIGHT AT LEAST 2 STANDARD DEVIATIONS (SD) BELOW THE MEAN HEIGHT FOR NORMAL CHILDREN OF THE SAME AGE AND GENDER. RENEWAL: GROWTH VELOCITY AND/OR TARGET HEIGHT.
Age Restrictions	
Prescriber Restrictions	PRESCRIBED BY OR GIVEN IN CONSULTATION WITH: ENDOCRINOLOGIST.
Coverage Duration	12 MONTHS
Other Criteria	RENEWAL: PHYSICIAN ATTESTATION OF IMPROVEMENT (I.E. INCREASED HEIGHT OR INCREASED GROWTH VELOCITY). THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION.

SOMATROPIN-NUTROPIN AND NUTROPIN AQ

Products Affected

- Nutropin AQ Nuspin

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	ATHLETIC ENHANCEMENT, ANTI-AGING PURPOSES, GROWTH FAILURE DUE TO CKD IF PATIENT HAS HAD A RENAL TRANSPLANT, OR GROWTH FAILURE WITH CLOSED EPIPHYSES.
Required Medical Information	INDUCTION - PATIENT'S HEIGHT AT LEAST 2 STANDARD DEVIATIONS (SD) BELOW THE MEAN HEIGHT FOR NORMAL CHILDREN OF THE SAME AGE AND GENDER. RENEWAL: GROWTH VELOCITY AND/OR TARGET HEIGHT.
Age Restrictions	
Prescriber Restrictions	PRESCRIBED BY OR GIVEN IN CONSULTATION WITH: ENDOCRINOLOGIST. FOR GROWTH HORMONE FAILURE DUE TO CRI: NEPHROLOGIST.
Coverage Duration	INITIAL AND RENEWAL: 12 MONTHS
Other Criteria	ALL DIAGNOSES EXCEPT FOR CHRONIC KIDNEY DISEASE (CKD): INITIAL: PREVIOUS TRIAL OF PREFERRED FORMULARY ALTERNATIVES NORDITROPIN AND GENOTROPIN PER FDA INDICATION. RENEWAL: PHYSICIAN ATTESTATION OF IMPROVEMENT (I.E. INCREASED HEIGHT OR INCREASED GROWTH VELOCITY). FOR GROWTH FAILURE SECONDARY TO CKD: PATIENT HAS NOT RECEIVED A RENAL TRANSPLANT. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION.

SONIDEGIB

Products Affected

- Odomzo

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	BASELINE SERUM CREATINE KINASE (CK) AND SERUM CREATININE LEVELS
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	

SORAFENIB TOSYLATE

Products Affected

- Nexavar

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	

SUNITINIB MALATE

Products Affected

- Sutent

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	GASTROINTESTINAL STROMAL TUMORS (GIST): TRIAL OF OR CONTRAINDICATION TO GLEEVEC.

TADALAFIL

Products Affected

- Cialis oral tablet 2.5 mg, 5 mg

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	ERECTILE DYSFUNCTION WITHOUT DIAGNOSIS OF BENIGN PROSTATIC HYPERPLASIA.
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	TRIAL OF ONE FORMULARY ALPHA BLOCKER SUCH AS DOXAZOSIN, TERAZOSIN, TAMSULOSIN OR ALFUZOSIN) AND ONE FORMULARY 5-ALPHA-REDUCTASE (SUCH AS FINASTERIDE OR DUTASTERIDE). APPLIES TO 2.5MG AND 5MG STRENGTHS ONLY. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION.

TALIMOGENE

Products Affected

- Imlygic injection suspension 10exp6 (1 million) PFU/mL, 10exp8 (100 million) PFU/mL

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	HISTORY OF PRIMARY OR ACQUIRED IMMUNODEFICIENT STATES, LEUKEMIA, LYMPHOMA, OR AIDS. PATIENT IS NOT CURRENTLY RECEIVING IMMUNOSUPPRESSIVE THERAPY.
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	IMLYGIC TO BE INJECTED INTO CUTANEOUS, SUBCUTANEOUS, AND OR NODAL LESIONS THAT ARE VISIBLE, PALPABLE, OR DETECTABLE BY ULTRASOUND GUIDANCE. NO CONCURRENT USE WITH PEMBROLIZUMAB, NIVOLUMAB, IPILIMUMAB, DABRAFENIB, TRAMETINIB, VEMURAFENIB, INTERLEUKIN-2, INTERFERON, DACARBAZINE, TEMOZOLOMIDE, PACLITAXEL, CARBOPLATIN, IMATINIB, MELPHALAN, IMIQUIMOD, OR RADIATION THERAPY.

TASIMELTEON

Products Affected

- Hetlioz

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	

TEDUGLUTIDE

Products Affected

- Gattex 30-Vial

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	18 YEARS OF AGE AND OLDER
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PATIENT IS DEPENDENT ON INTRAVENOUS PARENTERAL NUTRITION DEFINED AS REQUIRING PARENTERAL NUTRITION AT LEAST THREE TIMES PER WEEK.

TELOTRISTAT

Products Affected

- Xermelo

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	

TEMOZOLOMIDE

Products Affected

- Temodar intravenous

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	

TERIFLUNOMIDE

Products Affected

- Aubagio

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	

TERIPARATIDE

Products Affected

- Forteo

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	24 MONTHS OR MORE OF ANABOLIC THERAPY.
Required Medical Information	ONE OF THE FOLLOWING: (1) HIGH RISK FOR FRACTURES DEFINED AS ONE OF THE FOLLOWING: A) HISTORY OF OSTEOPOROTIC (I.E., FRAGILITY, LOW TRAUMA) FRACTURE(S). B) 2 OR MORE RISK FACTORS FOR FRACTURE (E.G., HISTORY OF MULTIPLE RECENT LOW TRAUMA FRACTURES, BMD T-SCORE LESS THAN OR EQUAL TO -2.5, CORTICOSTEROID USE, OR USE OF GNRH ANALOGS SUCH AS NAFARELIN, ETC.). C) NO PRIOR TREATMENT FOR OSTEOPOROSIS AND FRAX SCORE OF AT LEAST 20% FOR ANY MAJOR FRACTURE OR OF AT LEAST 3% FOR HIP FRACTURE. (2) UNABLE TO USE ORAL THERAPY (I.E., UPPER GASTROINTESTINAL PROBLEMS UNABLE TO TOLERATE ORAL MEDICATION, LOWER GASTROINTESTINAL PROBLEMS UNABLE TO ABSORB ORAL MEDICATIONS, TROUBLE REMEMBERING TO TAKE ORAL MEDICATIONS OR COORDINATING AN ORAL BISPHOSPHONATE WITH OTHER ORAL MEDICATIONS OR THEIR DAILY ROUTINE). (3) ADEQUATE TRIAL OF, INTOLERANCE TO, OR A CONTRAINDICATION TO BISPHOSPHONATES (E.G., ALENDRONATE, RISEDRONATE, IBANDRONATE).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	

TESTOSTERONE

Products Affected

- Androderm
- AndroGel transdermal gel in metered-dose pump 20.25 mg/1.25 gram (1.62 %)
- AndroGel transdermal gel in packet 1.62 % (20.25 mg/1.25 gram), 1.62 % (40.5 mg/2.5 gram)
- testosterone cypionate
- testosterone enanthate
- testosterone transdermal gel
- testosterone transdermal gel in packet

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D. ADDITIONAL CONSIDERATION FOR GENDER DYSPHORIA.
Exclusion Criteria	
Required Medical Information	MALE HYPOGONADISM CONFIRMED BY EITHER: 1) LAB CONFIRMED TOTAL SERUM TESTOSTERONE LEVEL OF LESS THAN 300 NG/DL OR 2) A LOW TOTAL SERUM TESTOSTERONE LEVEL AS INDICATED BY A LAB RESULT WITH A REFERENCE RANGE OBTAINED WITHIN 90 DAYS, OR 3) A FREE SERUM TESTOSTERONE LEVEL OF LESS THAN 5 PG/ML.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	LIFETIME OF MEMBERSHIP IN PLAN
Other Criteria	

TETRABENAZINE

Products Affected

- tetrabenazine

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	NEUROLOGIST
Coverage Duration	12 MONTHS
Other Criteria	

THALIDOMIDE

Products Affected

- Thalomid

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	

TOCILIZUMAB IV

Products Affected

- Actemra

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	RENEWAL: PHYSICIAN ATTESTATION OF IMPROVEMENT.
Age Restrictions	
Prescriber Restrictions	MODERATE TO SEVERE RHEUMATOID ARTHRITIS (RA)/POLYARTICULAR JUVENILE IDIOPATHIC ARTHRITIS (PJIA)/ SYSTEMIC JUVENILE IDIOPATHIC ARTHRITIS (SJIA): PRESCRIBED BY OR GIVEN IN CONSULTATION WITH A RHEUMATOLOGIST
Coverage Duration	INITIAL: RA: 6 MONTHS. PJIA: 5 MONTHS. SJIA: 12 MONTHS. RENEWAL: 12 MONTHS FOR ALL DIAGNOSES
Other Criteria	INITIAL: MODERATE TO SEVERE RA AND PJIA: PREVIOUS TRIAL OF HUMIRA AND ONE DMARD (DISEASE-MODIFYING ANTIRHEUMATIC DRUG) AGENT SUCH AS METHOTREXATE, LEFLUNOMIDE, HYDROXYCHLOROQUINE, OR SULFASALAZINE. INITIAL SJIA: PREVIOUS TRIAL WITH ONE DMARD (DISEASE-MODIFYING ANTIRHEUMATIC DRUG) AGENT SUCH AS METHOTREXATE, LEFLUNOMIDE, HYDROXYCHLOROQUINE, OR SULFASALAZINE.

TOCILIZUMAB SQ

Products Affected

- Actemra

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	RA RENEWAL: PHYSICIAN ATTESTATION OF IMPROVEMENT.
Age Restrictions	
Prescriber Restrictions	RHEUMATOID ARTHRITIS (RA): PRESCRIBED BY OR GIVEN IN CONSULTATION WITH A RHEUMATOLOGIST.
Coverage Duration	RA INITIAL: 6 MONTHS. RA RENEWAL: 12 MONTHS. GIANT CELL ARTERITIS: 12 MONTHS
Other Criteria	RA INITIAL : PREVIOUS TRIAL OF HUMIRA AND ONE DMARD (DISEASE-MODIFYING ANTIRHEUMATIC DRUG) AGENT SUCH AS METHOTREXATE, LEFLUNOMIDE, HYDROXYCHLOROQUINE, OR SULFASALAZINE.

TOFACITINIB

Products Affected

- Xeljanz
- Xeljanz XR

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	RENEWAL: PHYSICIAN ATTESTATION OF IMPROVEMENT.
Age Restrictions	
Prescriber Restrictions	RHEUMATOID ARTHRITIS: PRESCRIBED BY OR GIVEN IN CONSULTATION WITH A RHEUMATOLOGIST.
Coverage Duration	RA: INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.
Other Criteria	INITIAL: RHEUMATOID ARTHRITIS: PREVIOUS TRIAL OF HUMIRA AND ONE DMARD (DISEASE-MODIFYING ANTIRHEUMATIC DRUG) SUCH AS METHOTREXATE, LEFLUNOMIDE, HYDROXYCHLOROQUINE, OR SULFASALAZINE.

TOPICAL TRETINOIN

Products Affected

- tretinoin topical cream
- tretinoin topical gel 0.01 %, 0.025 %

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	WRINKLES, PHOTOAGING, MELASMA.
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	

TRABECTEDIN

Products Affected

- Yondelis

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	

TRAMETINIB DIMETHYL SULFOXIDE

Products Affected

- Mekinist oral tablet 0.5 mg, 2 mg

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	

TRASTUZUMAB

Products Affected

- Herceptin

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	BREAST CANCER, METASTATIC BREAST CANCER, GASTRIC CANCER: HER2 POSITIVE
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	B VS D COVERAGE CONSIDERATION.

TREPROSTINIL DIOLAMINE

Products Affected

- Orenitram

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	DOCUMENTED CONFIRMATORY PULMONARY ARTERIAL HYPERTENSION (PAH) DIAGNOSIS BASED ON RIGHT HEART CATHETERIZATION. PATIENT HAS NYHA-WHO FUNCTIONAL CLASS II-IV SYMPTOMS.
Age Restrictions	
Prescriber Restrictions	PRESCRIBED BY OR GIVEN IN CONSULTATION WITH A CARDIOLOGIST OR PULMONOLOGIST.
Coverage Duration	INITIAL AND RENEWAL: 12 MONTHS
Other Criteria	INITIAL: MEAN PULMONARY ARTERY PRESSURE (PAP) OF AT LEAST 25 MMHG OR GREATER, PULMONARY CAPILLARY WEDGE PRESSURE (PCWP) OF 15 MMHG OR LESS, PULMONARY VASCULAR RESISTANCE (PVR) GREATER THAN 3 WOOD UNITS. PATIENT DOES NOT HAVE SEVERE HEPATIC IMPAIRMENT. PREVIOUS OR CURRENT TREATMENT WITH ONE OF THE FOLLOWING: A FORMULARY PHOSPHODIESTERASE-5 INHIBITOR (E.G., SILDENAFIL [GENERIC FOR REVATIO] OR ADCIRCA [TADALAFIL]) OR AN ENDOTHELIN RECEPTOR ANTAGONIST (E.G., TRACLEER [BOSENTAN], LETAIRIS [AMBRISENTAN], OR OPSUMIT [MACITENTAN]). TRIAL OF A FORMULARY PHOSPHODIESTERASE-5 INHIBITOR OR ENDOTHELIN RECEPTOR ANTAGONIST IS NOT REQUIRED IF THE PATIENT WAS PREVIOUSLY STABLE ON ORENITRAM. RENEWAL: PATIENT SHOWS IMPROVEMENT FROM BASELINE IN THE 6-MINUTE WALK DISTANCE OR PATIENT HAS A STABLE 6-MINUTE WALK DISTANCE WITH A STABLE/ IMPROVED WHO FUNCTIONAL CLASS.

TREPROSTINIL INHALED

Products Affected

- Tyvaso

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	DOCUMENTED CONFIRMATORY PULMONARY ARTERIAL HYPERTENSION (PAH) DIAGNOSIS BASED ON RIGHT HEART CATHETERIZATION. PATIENT HAS NYHA-WHO FUNCTIONAL CLASS III-IV SYMPTOMS.
Age Restrictions	
Prescriber Restrictions	PRESCRIBED BY OR IN CONSULTATION WITH A CARDIOLOGIST OR PULMONOLOGIST
Coverage Duration	INITIAL AND RENEWAL: 12 MONTHS
Other Criteria	THIS DRUG MAYBE COVERED UNDER MEDICARE PART B OR D DEPENDING UPON THE CIRCUMSTANCES. INFORMATION MAY NEED TO BE SUBMITTED DESCRIBING THE USE AND SETTING OF THE DRUG TO MAKE THE DETERMINATION. NEBULIZER THERAPY IS COVERED UNDER PART B FOR PATIENTS WHO ARE USING THE MEDICATION VIA A NEBULIZER IN THEIR OWN HOME. THOSE WHO ARE NOT USING IT IN THEIR HOME WILL BE COVERED UNDER PART D. INITIAL: MEAN PULMONARY ARTERY PRESSURE (PAP) OF AT LEAST 25 MMHG OR GREATER, PULMONARY CAPILLARY WEDGE PRESSURE (PCWP) OF 15 MMHG OR LESS, PULMONARY VASCULAR RESISTANCE (PVR) GREATER THAN 3 WOOD UNITS. RENEWAL: PATIENT SHOW IMPROVEMENT FROM BASELINE IN THE 6-MINUTE WALK DISTANCE OR PATIENT HAS A STABLE 6-MINUTE WALK DISTANCE WITH A STABLE/ IMPROVED WHO FUNCTIONAL CLASS.

TREPROSTINIL SODIUM INJECTABLE

Products Affected

- Remodulin

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	COVERED UNDER LOCAL COVERAGE POLICY OF APPLICABLE MEDICARE DMERC.
Required Medical Information	FORMULARY DRUG ADMINISTERED IN A LONG TERM CARE FACILITY TO A PATIENT WHOSE PART A COVERAGE HAS EXPIRED OR FORMULARY DRUG NOT ADMINISTERED VIA AN IMPLANTABLE PUMP OR AN EXTERNAL PUMP OR DRUG ADMINISTERED VIA AN IMPLANTABLE PUMP/AN EXTERNAL PUMP. DOCUMENTED CONFIRMATORY PULMONARY ARTERIAL HYPERTENSION (PAH) DIAGNOSIS BASED ON RIGHT HEART CATHETERIZATION.
Age Restrictions	
Prescriber Restrictions	PRESCRIBED BY OR IN CONSULTATION WITH A CARDIOLOGIST OR PULMONOLOGIST
Coverage Duration	INITIAL AND RENEWAL: 12 MONTHS

PA Criteria	Criteria Details
Other Criteria	<p>INITIAL: MEAN PULMONARY ARTERY PRESSURE (PAP) OF AT LEAST 25 MMHG OR GREATER, PULMONARY CAPILLARY WEDGE PRESSURE (PCWP) OF 15 MMHG OR LESS, PULMONARY VASCULAR RESISTANCE (PVR) GREATER THAN 3 WOOD UNITS. CONTINUATION OF CURRENT REMODULIN THERAPY: PATIENT MUST HAVE NYHA/WHO FC II-IV SYMPTOMS. NEW REQUESTS FOR REMODULIN THERAPY: PATIENT MUST HAVE NYHA/WHO FC III-IV SYMPTOMS. NEW REQUESTS FOR REMODULIN THERAPY FOR PATIENTS WITH NYHA/WHO FC II SYMPTOMS REQUIRES A TRIAL OF OR CONTRAINDICATION TO A PHOSPHODIESTERASE-5 INHIBITOR (PDE-5) (E.G., REVATIO, ADCIRCA) OR AN ENDOTHELIN RECEPTOR ANTAGONIST (ERA) (E.G., LETAIRIS, OPSUMIT, TRACLEER). RENEWAL: PATIENT SHOW IMPROVEMENT FROM BASELINE IN THE 6-MINUTE WALK DISTANCE OR PATIENT HAS A STABLE 6-MINUTE WALK DISTANCE WITH A STABLE/ IMPROVED WHO FUNCTIONAL CLASS.</p>

TRIENTINE

Products Affected

- Syprine

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	KNOWN FAMILY HISTORY OF WILSON'S DISEASE OR PHYSICAL EXAMINATION CONSISTENT WITH WILSON'S DISEASE. PLASMA COPPER-PROTEIN CERULOPLASMIN LESS THAN 20 MG/DL. LIVER BIOPSY POSITIVE FOR AN ABNORMALLY HIGH CONCENTRATION OF COPPER (GREATER THAN 250 MCG/G DRY WEIGHT) OR THE PRESENCE OF KAYSER-FLEISCHER RINGS. MAINTENANCE OF A REDUCED COPPER DIETARY INTAKE (LESS THAN 2 MG COPPER PER DAY).
Age Restrictions	
Prescriber Restrictions	PRESCRIBED BY OR GIVEN IN CONSULTATION WITH A HEPATOLOGIST.
Coverage Duration	12 MONTHS
Other Criteria	PREVIOUS TRIAL OF OR CONTRAINDICATION TO PENICILLAMINE (DEPEN).

TRIFLURIDINE/TIPIRACIL

Products Affected

- Lonsurf oral tablet 15-6.14 mg, 20-8.19 mg

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	

USTEKINUMAB

Products Affected

- Stelara subcutaneous syringe

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	INITIAL: PLAQUE PSORIASIS: MODERATE TO SEVERE PLAQUE PSORIASIS INVOLVING GREATER THAN OR EQUAL TO 5% BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, OR GENITAL AREA. RENEWAL FOR PSORIATIC ARTHRITIS OR PLAQUE PSORIASIS: PHYSICIAN ATTESTATION OF IMPROVEMENT.
Age Restrictions	
Prescriber Restrictions	PRESCRIBED BY OR GIVEN IN CONSULTATION WITH: PSORIATIC ARTHRITIS: DERMATOLOGIST OR RHEUMATOLOGIST. PLAQUE PSORIASIS: DERMATOLOGIST. CROHN'S DISEASE: GASTROENTEROLOGIST.
Coverage Duration	INITIAL: PSA, PSO, CD: 4 MONTHS. CD WITH PREVIOUS DOSE IV: 2 MONTHS. RENEW ALL: 12 MO
Other Criteria	INITIAL: PSORIATIC ARTHRITIS (PSA): PREVIOUS TRIAL OF HUMIRA FOLLOWED BY ONE OF THE FOLLOWING PREFERRED AGENTS: CIMZIA, OTEZLA, OR COSENTYX. PLAQUE PSORIASIS (PSO): PREVIOUS TRIAL OF HUMIRA FOLLOWED BY ONE OF THE FOLLOWING PREFERRED AGENTS: COSENTYX OR OTEZLA. CROHN'S DISEASE (CD): PREVIOUS TRIAL OF OR CONTRAINDICATION TO PREFERRED TNF INHIBITORS: HUMIRA FOLLOWED BY CIMZIA.

USTEKINUMAB IV

Products Affected

- Stelara intravenous

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	PRESCRIBED BY OR GIVEN IN CONSULTATION WITH A GASTROENTEROLOGIST.
Coverage Duration	2 MONTHS
Other Criteria	PREVIOUS TRIAL OF OR CONTRAINDICATION TO THE PREFERRED TNF INHIBITORS: HUMIRA FOLLOWED BY CIMZIA. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.

VALBENZAZINE TOSYLATE

Products Affected

- Ingrezza

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	PATIENT HAS BEEN USING ANTIPSYCHOTIC MEDICATIONS OR METOCLOPRAMIDE FOR AT LEAST 3 MONTHS (OR AT LEAST 1 MONTH IF PATIENT IS 60 YEARS OF AGE OR OLDER) PER PHYSICIAN ATTESTATION
Age Restrictions	
Prescriber Restrictions	PRESCRIBED BY OR GIVEN IN CONSULTATION WITH A NEUROLOGIST OR MOVEMENT DISORDER SPECIALIST
Coverage Duration	12 MONTHS
Other Criteria	MODERATE TO SEVERE TARDIVE DYSKINESIA HAS BEEN PRESENT FOR AT LEAST 3 MONTHS

VANDETANIB

Products Affected

- Caprelsa oral tablet 100 mg, 300 mg

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	

VEMURAFENIB

Products Affected

- Zelboraf

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	BRAFV600E MUTATION
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	

VENETOCLAX

Products Affected

- Venclexta oral tablet 10 mg, 100 mg, 50 mg
- Venclexta Starting Pack

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	

VINCRIStINE SULFATE LIPOSOMAL

Products Affected

- Marqibo

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.

VISMODEGIB

Products Affected

- Erivedge

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	

ZIV-AFLIBERCEPT

Products Affected

- Zaltrap

PA Criteria	Criteria Details
Covered Uses	ALL FDA APPROVED INDICATIONS NOT OTHERWISE EXCLUDED FROM PART D.
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	

INDEX

Actemra.....	247, 248	butalbital-acetaminophen oral tablet 50-325 mg.....	103
Acthar H.P.....	45	butalbital-acetaminophen-caff oral capsule 50-325-40 mg.....	103
Adcirca.....	189	butalbital-acetaminophen-caff oral tablet 50-325-40 mg.....	103
Adempas.....	209	butalbital-aspirin-caffeine oral capsule...	103
Afinitor Disperz.....	88	Cabometyx oral tablet 20 mg, 40 mg, 60 mg.....	36
Afinitor oral tablet 10 mg, 2.5 mg, 5 mg, 7.5 mg.....	88	Capacet.....	103
Alecensa.....	8	Caprelsa oral tablet 100 mg, 300 mg.....	263
Alunbrig.....	32	carisoprodol.....	110
amitriptyline.....	120	Cerdelga.....	73
amitriptyline-chlordiazepoxide.....	120	chlorzoxazone.....	110
amoxapine.....	120	Cialis oral tablet 2.5 mg, 5 mg.....	236
Ampyra.....	51	Cimzia.....	42
Anadrol-50.....	182	Cimzia Powder for Reconst.....	42
Androderm.....	244	Cinqair.....	206
AndroGel transdermal gel in metered-dose pump 20.25 mg/1.25 gram (1.62 %).	244	Cinryze.....	34
AndroGel transdermal gel in packet 1.62 % (20.25 mg/1.25 gram), 1.62 % (40.5 mg/2.5 gram).....	244	clomipramine.....	120
armodafinil.....	160	CombiPatch.....	106
Ascomp with Codeine.....	103	Cometriq.....	35
Aubagio.....	242	Copaxone subcutaneous syringe 40 mg/mL.....	96
Austedo oral tablet 12 mg, 6 mg, 9 mg.....	61	Corlanor.....	137
Avastin.....	26	Cosentyx (2 Syringes).....	220
Avonex (with albumin).....	134	Cosentyx Pen (2 Pens).....	220
Avonex intramuscular pen injector kit...	134	Cotellic.....	44
Avonex intramuscular syringe kit.....	134	Cuprimine.....	193
Bavencio.....	19	cyclobenzaprine oral tablet.....	110
Beleodaq.....	24	cyproheptadine.....	111
Bendecka.....	25	Cyramza.....	204
Benlysta intravenous.....	23	Daklinza.....	48
benztropine.....	100	Daraprim.....	201
Betaseron subcutaneous kit.....	135	Darzalex.....	52
bexarotene.....	27	deferoxamine.....	56
Blinicyto intravenous kit.....	28	Depen Titratabs.....	194
Bosulif oral tablet 100 mg, 500 mg.....	30	desipramine.....	120
Botox injection recon soln 100 unit, 200 unit.....	31	diclofenac sodium topical gel 3 %.....	64
buprenorphine HCl sublingual.....	180	Digitek oral tablet 125 mcg, 250 mcg.....	105
Butalbital Compound W/Codeine.....	103	DIGOXIN 0.25 MG/ML SYRINGE.....	105
butalbital-acetaminop-caf-cod.....	103	digoxin injection solution.....	105
		digoxin oral solution 50 mcg/mL.....	105

digoxin oral tablet 125 mcg, 250 mcg.....	105	Gilenya.....	94
diphenhydramine HCl oral elixir.....	112	Gilotrif.....	7
diphenoxylate-atropine.....	113	Glatopa.....	96
doxepin oral.....	120	glyburide.....	108
dronabinol.....	38	glyburide micronized.....	108
Duavee.....	106	glyburide-metformin.....	108
Duexis.....	125	guanfacine oral tablet.....	104
Dupixent.....	68	Haegarda.....	34
Emflaza oral suspension.....	57	Harvoni.....	142
Emflaza oral tablet 18 mg, 30 mg, 36 mg, 6 mg.....	57	Herceptin.....	253
Empliciti.....	75	Hetlioz.....	238
Enbrel.....	85	Humatrope.....	228
Enbrel SureClick.....	85	Humira.....	5
Epclusa.....	227	Humira Pediatric Crohn's Start.....	5
EPOGEN 10,000 UNITS/ML VIAL SDV, P/F, OUTER.....	81	Humira Pen.....	5
Epogen injection solution 2,000 unit/mL, 20,000 unit/2 mL, 20,000 unit/mL, 3,000 unit/mL, 4,000 unit/mL.....	81	Humira Pen Crohn's-UC-HS Start.....	5
epoprostenol (glycine).....	79	Humira Pen Psoriasis-Uveitis.....	5
Erivedge.....	267	hydroxyprogesterone caproate.....	123
Esbriet oral capsule.....	197	hydroxyzine HCl intramuscular.....	114
Esbriet oral tablet 267 mg, 801 mg.....	197	hydroxyzine HCl oral solution 10 mg/5 mL.....	114
estazolam oral tablet 1 mg, 2 mg.....	121	hydroxyzine HCl oral tablet.....	114
estradiol oral.....	106	hydroxyzine pamoate.....	114
estradiol transdermal patch semiweekly..	106	Ibrance.....	183
estradiol transdermal patch weekly.....	106	Iclusig oral tablet 15 mg, 45 mg.....	199
estradiol-norethindrone acet.....	106	Ilaris (PF).....	37
estropipate.....	106	imatinib oral tablet 100 mg, 400 mg.....	127
eszopiclone.....	109	Imbruvica.....	124
Exjade.....	54	Imfinzi.....	69
Exondys 51.....	87	imipramine HCl.....	120
Extavia subcutaneous kit.....	135	imipramine pamoate.....	120
Farydak.....	186	imiquimod.....	128
fentanyl citrate.....	93	Imlygic injection suspension 10exp6 (1 million) PFU/mL, 10exp8 (100 million) PFU/mL.....	237
Ferriprox.....	55	Impavido.....	156
Flector.....	63	indomethacin oral capsule 25 mg, 50 mg..	115
flurazepam oral capsule 15 mg, 30 mg.....	121	indomethacin oral capsule, extended release.....	115
Forteo.....	243	Inflectra.....	131
Fyavolv oral tablet 1-5 mg-mcg.....	106	Ingrezza.....	262
Gattex 30-Vial.....	239	Inlyta oral tablet 1 mg, 5 mg.....	20
Gazyva.....	170	Intron A injection.....	133
Genotropin.....	231	Iprivask.....	60
Genotropin MiniQuick.....	231	Iressa.....	95

Jadenu.....	54	Mimvey Lo.....	106
Jadenu Sprinkle.....	54	Mircera injection syringe 100 mcg/0.3 mL, 200 mcg/0.3 mL, 50 mcg/0.3 mL, 75 mcg/0.3 mL.....	84
Jakafi.....	215	Myobloc intramuscular solution 10,000 unit/2 mL, 2,500 unit/0.5 mL, 5,000 unit/mL.....	31
Jinteli.....	106	Natpara.....	187
Juxtapid oral capsule 10 mg, 20 mg, 30 mg, 40 mg, 5 mg, 60 mg.....	147	Nexavar.....	234
Kalydeco.....	138, 139	nifedipine oral capsule.....	104
Kanuma.....	218	Ninlaro.....	140
ketorolac oral.....	116	nitrofurantoin.....	102
Keveyis.....	62	nitrofurantoin macrocrystal.....	102
Kevzara.....	217	nitrofurantoin monohyd/m-cryst.....	102
Keytruda intravenous recon soln.....	192	Norditropin FlexPro.....	231
Keytruda intravenous solution.....	192	norethindrone ac-eth estradiol oral tablet 0.5-2.5 mg-mcg, 1-5 mg-mcg.....	106
Kineret.....	12	Northera.....	67
Kisqali.....	207	nortriptyline.....	120
Kisqali Femara Co-Pack oral tablet 200 mg/day(200 mg x 1)-2.5 mg, 400 mg/day(200 mg x 2)-2.5 mg, 600 mg/day(200 mg x 3)-2.5 mg.....	207	Nplate.....	213
Korlym.....	155	Nucala.....	151
Kynamro.....	157	Nuplazid.....	196
Kyprolis.....	40	Nutropin AQ Nuspin.....	232
Lanoxin oral tablet 187.5 mcg, 62.5 mcg.....	105	Ocaliva.....	169
Lartruvo.....	173	Ocrevus.....	171
Lazanda.....	92	Odomzo.....	233
Lemtrada.....	9	Ofev.....	165
Lenvima.....	144	Olysio.....	223
Letairis.....	77	Omnitrope.....	228
lidocaine topical adhesive patch,medicated.....	145	Oncaspar.....	17
lidocaine topical ointment.....	145	Onfi oral suspension.....	43
Lonsurf oral tablet 15-6.14 mg, 20-8.19 mg.....	259	Onfi oral tablet 10 mg, 20 mg.....	43
Lopreeza.....	106	Opdivo.....	168
Lynparza.....	172	Opsumit.....	77
Marqibo.....	266	Orencia.....	3
meclizine oral tablet 12.5 mg, 25 mg.....	117	Orencia (with maltose).....	2
megestrol oral suspension 400 mg/10 mL (40 mg/mL).....	118	Orencia ClickJect.....	3
megestrol oral tablet.....	118	Orenitram.....	254
Mekinist oral tablet 0.5 mg, 2 mg.....	252	Orfadin.....	167
Menest oral tablet 0.3 mg, 0.625 mg, 1.25 mg.....	106	Orkambi.....	150
methocarbamol oral.....	110	Otezla.....	13
Mimvey.....	106	Otezla Starter.....	13
		paroxetine HCl oral tablet.....	119
		paroxetine HCl oral tablet extended release 24 hr.....	119

Paxil oral suspension.....	119	Saizen click.easy.....	228
Pennsaid topical solution in metered-dose pump.....	64	Serostim subcutaneous recon soln 4 mg, 5 mg, 6 mg.....	229
Perjeta.....	195	sildenafil intravenous.....	190
perphenazine-amitriptyline.....	120	sildenafil oral.....	189
Phenadoz.....	101	Siliq.....	33
phenobarbital.....	122	Simponi.....	99
Plegridy.....	134	Simponi ARIA.....	98
Pomalyst.....	198	Sirturo.....	22
Portrazza.....	163	Sovaldi.....	225
Praluent Pen.....	10	Sprycel oral tablet 100 mg, 140 mg, 20 mg, 50 mg, 70 mg, 80 mg.....	53
Premarin oral.....	106	Stelara intravenous.....	261
Premphase.....	106	Stelara subcutaneous syringe.....	260
Prempro.....	106	Stivarga.....	205
Procrit injection solution 10,000 unit/mL, 2,000 unit/mL, 20,000 unit/2 mL, 20,000 unit/mL, 3,000 unit/mL, 4,000 unit/mL, 40,000 unit/mL.....	81	Strensiq.....	14
Promacta.....	76	Surmontil.....	120
promethazine injection solution.....	101	Sutent.....	235
promethazine oral.....	101	Sylatron.....	191
promethazine rectal.....	101	Sylvant.....	222
Promethegan.....	101	SymlinPen 120.....	200
protriptyline.....	120	SymlinPen 60.....	200
quinine sulfate.....	203	Synagis.....	184
Radicava.....	70	Syndros.....	39
Ravicti.....	97	Synribo.....	174
Rebif (with albumin).....	134	Syprine.....	258
Rebif Rebidose.....	134	Tafinlar.....	47
Rebif Titration Pack.....	134	Tagrisso.....	181
Regranex.....	21	Taltz Autoinjector.....	141
Relistor oral.....	153	Taltz Syringe.....	141
Relistor subcutaneous solution.....	152	Tarceva oral tablet 100 mg, 150 mg, 25 mg.....	80
Relistor subcutaneous syringe.....	152	Targretin topical.....	27
Remicade.....	129	Tasigna.....	164
Remodulin.....	256	Tecentriq.....	18
Repatha Pushttronex.....	89	Tecfidera oral capsule, delayed release(DR/EC) 120 mg, 120 mg (14)- 240 mg (46), 240 mg.....	65
Repatha SureClick.....	89	Technivie.....	176
Repatha Syringe.....	89	temazepam oral capsule 15 mg, 30 mg....	121
Revlimid.....	143	Temodar intravenous.....	241
Rituxan.....	211	Tencon oral tablet 50-325 mg.....	103
Rituxan Hycela.....	212	testosterone cypionate.....	244
Rubraca.....	214	testosterone enanthate.....	244
Rydapt.....	154	testosterone transdermal gel.....	244
Saizen.....	228		

testosterone transdermal gel in packet	244	Zinbryta.....	50
tetrabenazine.....	245	zolpidem oral.....	109
Thalomid.....	246	Zomacton.....	228
Tracleer.....	77	Zorbtive.....	230
tretinoin topical cream.....	250	Zydelig.....	126
tretinoin topical gel 0.01 %, 0.025 %.....	250	Zykadia.....	41
triazolam oral tablet 0.125 mg, 0.25 mg..	121	Zytiga.....	4
trihexyphenidyl.....	100		
trimipramine.....	120		
Tymlos.....	1		
Tysabri.....	162		
Tyvaso.....	255		
Unituxin.....	66		
Uptravi oral tablet 1,000 mcg, 1,200 mcg, 1,400 mcg, 1,600 mcg, 200 mcg, 400 mcg, 600 mcg, 800 mcg.....	221		
Uptravi oral tablets,dose pack.....	221		
Vectibix.....	185		
Velcade.....	29		
Venclexta oral tablet 10 mg, 100 mg, 50 mg.....	265		
Venclexta Starting Pack.....	265		
Viekira Pak.....	178		
Viekira XR.....	178		
Vimizim.....	74		
Vimovo.....	161		
Votrient.....	188		
Xadago.....	216		
Xalkori.....	46		
Xeljanz.....	249		
Xeljanz XR.....	249		
Xermelo.....	240		
Xgeva.....	59		
Xifaxan oral tablet 200 mg, 550 mg.....	208		
Xiidra.....	146		
Xolair.....	175		
Xtandi.....	78		
Yervoy.....	136		
Yondelis.....	251		
zaleplon.....	109		
Zaltrap.....	268		
Zebutal oral capsule 50-325-40 mg.....	103		
Zejula.....	166		
Zelboraf.....	264		
Zepatier.....	71		